Journal of Agriculture and Forest Meteorology Research

JAFMR, 4(1): 336-339 www.scitcentral.com ISSN: 2642-0449

Review Article: Open Access

Previous Processing – Sure Way for Promotion of Microbiological Safety of Germinated Broad Bean Seeds

Gurbanov NH*, Gadimova NS, Akhundova NA, Mageramova MH and Gurbanova RI

*Department of Food Products Technology, Azerbaijan State University of Economics (UNEC), Baku, Azerbaijan.

Received August 22, 2019; Accepted September 11, 2019; Published January 08, 2020

ABSTRACT

The microbiological contamination of broad beans seeds in the process of germination in husked and natural form was studied. It was shown that preliminary short-term hydrothermal treatment and removal of the fruit coat of broad bean seeds provides an increase in the dynamics of their germination and microbiological stability in the process of germination. Therefore, this method of germination of broad bean seeds can be recommended for practical use.

Keywords: Broad bean seeds, Germination in peeled form, Microbiological indices

INTRODUCTION

Germinated cereal products, such as natural are being used in health-improving diets [1-3]. Research conducted last years in this direction for the inventions of the products of functional purpose have obtained huge scales [4-9].

Based on it, our research with broad bean seeds, which are called "pakhla" in the folklore like traditional proteinbearing raw material, widely used in cooking, in order to get germinated composition and food products from its, deserves attention [10-14].

Proposed modified method by us of deriving the germinated broad bean seeds, in our view, replies above indicated specifications, makes impossible promotion of the level of microbial contamination of the germinated production. As in the process of germination of the latter twice – at the start of preparation of the seeds and their preliminary soaking results in significant decrease of microbiological filth of the raw material, due to smooth regulation of the preliminary short hydrothermal treatment for removing the most infected details of the beans – seed membranes. With that, the process of preparations of the germinated samples, promotion of their appearance and improvement of nutritive value of the ready production increases.

Thus, in order to decrease the microbial contamination of germinated seeds of broad beans and the time of its germination, increase the activity of endoenzymes, increase the receiver of the germinant in a developed process, shortlived special wet-heat treatment and removing bran covering, before soaking the raw material, was applied. After that the peeled seeds (seed lobe) were soaked in drinkable municipal water at the temperature of $20^{\circ} \pm 4^{\circ}$ C (hydro modulus composed 1:10) during 21 h, then were germinated during 72 h. Every 12 h the amounts of the germinated seeds to their total number were written down (Figure 1).

Corresponding author: Gurbanov NH, Assistant Professor, Department of Food Products Technology, Azerbaijan State University of Economics (UNEC), AZ-1134, Baku City, 4th District, Asi Maharramov Street 68, Apartment 86, Baku, Azerbaijan, Tel: +994 050 338 77 20; E-mail: nusratgurbanov@hotmail.com

Citation: Gurbanov NH, Gadimova NS, Akhundova NA, Mageramova MH & Gurbanova RI. (2021) Previous Processing – Sure Way for Promotion of Microbiological Safety of Germinated Broad Bean Seeds. J Agric Forest Meteorol Res, 4(1): 336-339.

Copyright: ©2020 Gurbanov NH, Gadimova NS, Akhundova NA, Mageramova MH & Gurbanova RI. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1. The dynamics of sprouting of the broad bean seeds depending on germination method.

Traditional method (natural unpeeled seeds) - Suggested method (peeled seeds)

It should be noticed, that in the process of germination of peeled bean samples by suggested method, they produce

double sprouts (Figure 2). However, at natural unpeeled samples it is hardly in evidence, in other words they possess single sprouts. Besides, the process of germination happens much more slowly in comparison to traditional way of germination.



Figure 2. External appearance of the seeds of broad beans before and after germination depending on way of preproduction of the raw material (A) natural seeds; (B) Peeled seeds; (C) Germinated by traditional way; (D) Germinated by suggested way.

RESULTS AND DISCUSSION

The results of research alternations of morphological factors and appearance of the seeds in the process of germination are shown in the **Table 1**.

J Agric Forest Meteorol Res, 4(1): 336-339

Gurbanov NH, Gadimova NS, Akhundova NA, Mageramova MH & Gurbanova RI

Kind, used for germination of	Average dimensions of dry bean seeds, sm		Middle dimensions of alone and binominal sprouts after germination, sm		Middle mass of unpeeled and peeled germinated seeds with germinant (g)		Alternation of mass of the seeds after germination (%)	
secus	Length	Width	1 st	2 nd	Before germination	After germination	ger mination, (70)	
With a husk (unpeeled)	3.0	2.0	1.9	-	2.7	7.3	267	
Without a husk (peeled)	2.8	1.8	2.5	1.6	3.0	11.5	380	

Table 1. Alternation of morphological factors and appearance of the broad bean seeds in the process of germination.

As seen from **Table 1 and Figure 1**, preproduction and removal bran covering from bean seeds, at the time as improvement of appearance of sprouts increase the dynamic of their germination. The microbial contamination of broad bean seeds, germinated by traditional and suggested way, was determined by well-knowing methods [15-17].

available in the **Table 2**. As seen as from the factors in the **Table 2**, natural seeds in the process of preparation for germination possess twice as large contamination, than the seeds removed from bran covering by applying preliminary wet-heat treatment. An analogous scene is observable in the ready germinated samples.

The determination of alternation results of microbiological factors of broad bean seeds in the process of germination is

Table 2. Microbiological attributes unpeeled and peeled broad bean seeds in the process of germination.

No.	The technology of	Microbiological attributes afloat seeds, CFU, g							
	germination	Mesophilic aerobic and facultative anaerobic microorganisms	CB in 0.1 g seeds	Yeast	Fungus				
Ι	Germination of natural unpeeled seeds								
	Seeds after soaking	$6.0 imes 10^5$	Is absent	Not detected	Not detected				
	Seeds after germination	$1.6 \times \cdot 10^{12}$	Elicited	$3.0 \times \cdot 10^{3}$	30				
II	Germination of peeled seeds								
	Seed lobes after soaking	$3.0 \times \cdot 10^{2}$	Is absent	Not detected	Not detected				
	Seed lobes after germination	$2.5 \times \cdot 10^{6}$	Is absent	<15	<15				

Taking into account the high-quality and quantitative factors of germinated broad bean seeds by suggested technology and perspective of their usage for deriving functional nutritional products, it is considered to be desirable the additional research of the condition of microbial contamination of germinated samples in the process of storage. Being that, given type of the production in spite of popularity, possess short term of storage. According to the information shown in the **Table 2**, the appliance of suggested technology of broad bean seeds germination by usage of peeled seed lobes under soaking, stipulates their reduction of contamination about 2 h, which further reflects the deplumation security layer of germinated germinant, which composes about $2.5 \times \cdot 10^6$ (of CFU/g), which is close to the ratio for ready production ($5.0 \times \cdot 10^5$ CFU/g for cereals). All of this confirms practicability of

J Agric Forest Meteorol Res, 4(1): 336-339

broad bean seeds germination after their husking via use of short-time wet-heat treatment.

For this reason, in order to increase the drawdown period of germinant (including seed lobe) made from broad bean seeds and for prevention of their recontamination with microorganisms, germinated samples after well washing were dried on the air, kept in polythene bag weighing of 100-150 g in the refrigerator at $+6^{\circ}$ C till 5 days. Researches have shown that, safety preservation of germinated seeds in polythene bags is provided maximum till 4 days, as after 5 days of their preservation in the bags the bacterial load growths above accepted level of safety [18].

The research allow to draw the conclusion that, the usage of directed regime of bean seeds germination in peeled form allows not only to speed up the time for germination, but also to derive seed mass with the least microbial contamination.

CONCLUSION

- 1. Short-term moisture-thermal treatment of broad bean seeds with the aim of peeling and reducing antinutritional factors in them, provides a simultaneous improvement in seed yield and their microbiological parameters during germination.
- 2. Removing the fruit coat of broad bean seeds by means of thermal treatment approximately halves the microbiological contamination of the original seeds.

REFERENCES

- 1. Polyakova SP, Sidorova OA (2012) Improving stability of confectionary and bakery production to bacteriological damage. Food Industry 5: 16-18.
- 2. Pogojeva AV (2006) Valuation of utilization efficiency of bread, made from germinated cereal, in dietic therapy late adulthood with cardiovascular diseases. Problems Nutr 5: 45-48.
- Ehrenbergerova J (2008) Different barley cultivars as a source of green mass for improving nutrient balance in human diet. Fortschritte in der Saatguttechnologie und – untersuchung – ertragsorientierte. Züchtungsstrategien für neue Verwertungsmöglichkeiten, pp: 91-94.
- 4. Sabelnikova EA, Parfenov AI, Krums LM, Belousova NL (2010) The problem of the organization of dietary nutrition for patients with gluten-sensitive celiac disease in Russia. Exp Clin Gastroenterol 3: 107-110.
- 5. Bogatiryeva TG, Labutina NV, Belyavskaya IG, Yudina TA (2016) The technology of rye-wheaten bread in terms of corn ferments. Bread Products 9: 49-51.
- 6. Veber AL, Kazidub NG, Leonova SA, Jiarno M (2017) The deriving of biologically active element from germinated bean by way of its future reference. Bread Products 6: 35-39.

- Vidal-Valverde C, Frias J, Sotomayor C, Diaz-Pollan C, Fernandez M, et al. (1998) Nutrients and anti-nutritional factors in Faba beans as affected by processing. Z Lebensm Unters Forsch A 207: 140-145.
- Borisenko LA, Bratsikhin AA, Borisenko AA (2009) New types of cereal based meat semi-finished products. Food Industry 10: 16-17.
- 9. Leonova SA, Nutredinova OF, Fazilov MZ (2015) Technology of obtaining a national cereal product from the germinated grain of oats with addition of apples. Bread Products 9: 52-53.
- Gurbanov NH, Gurbanova RI, Maharramova MH, Isgenderova MM (2017) Combined proteinic-vitaminic salad in terms of germinated broad bean seeds. Collection of news «National academy of sciences of Azerbaijan. Ganja Department 4: 134-144.
- 11. Gurbanov NH, Gurbanova RI, Maharramova MH, Isgenderova MM (2017) Variance examination of general chemical and amino acid composition, ferments strength of peeled broad bean seeds in the process of germination. Proceedings of Azerbaijan Agrarian University, Ganja 3: 45-48.
- Gurbanov NH, Omarova EM, Gurbanova RI, Yusifzade SN (2014) Grain milk from germinated broad bean seeds for production of combined fermented milk drinks of functional purpose. Conference for 13th Anniversary of the National Institute of Food Technology, KYIV, pp: 56-57.
- Gurbanov NH, Omraova EM, Gasanova ZP (2015) Quantitative study of amino acid composition of faba bean seeds before and after of bioactivation. International Scientific and Technical Conference, Mogilyov, p: 148.
- Gurbanov NH, Maharramova MH, Gurbanova RI (2014) Bio activation of broad beans (Faba vulgaris Moench) for deriving food compositions purpose of use. Microbiology Institute Transactions of Azerbaijan National Academy of sciences, Baku 1: 149-154.
- 15. GOST 26669-85 «Nutritive and flavor products. Preparations of samples for microbiology testing».
- 16. GOST 10444.15-94 «Nutritive products. Methods of quantity measurement of mesophilic, aerobic and facultative anaerobic microorganisms».
- 17. GOST P 54005-2010 «Nutritive products. Methods of detecting and quantity determination of bacteriums of the family Enterobacteriaceae».
- 18. Alekhina NN (2017) Investigation of the influence of the hop structure on microbiological indicators of the bread quality from bio activated wheat grain. Bread Products 6: 60-61.