

Plant Responses to Heat Stress and Thermotolerance

A Hemantaranjan^{1*}, A Nishant Bhanu² and Shivani Lalotra¹

¹Department of Plant Physiology, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi - 221005, India

²Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi - 221005, India.

Received September 09, 2019; Accepted November 06, 2019; Published January 06, 2020

ABSTRACT

With increasing challenges posed by climate change, it is predicted that warming, drought, floods and storm events will become even more frequent and severe and will further reduce crop yields, especially in the tropics and subtropics. Temperatures above the normal optimum cause heat stress (HS) at different levels in all living organisms. Heat stress disturbs cellular homeostasis and causes denaturation and dysfunction in numerous proteins, leading to severe retardation in growth, development and even death. In plants, the major sites of heat stress injury are the oxygen-evolving complex (OEC) along with linked biochemical reactions in photosystem II (PSII). Perception of heat stress by plants usually triggers sensors at the plasma membrane and causes a transient opening of Ca^{2+} channels, possibly via modulation of membrane fluidity. Heat stress differentially influences the constancy of various proteins, membranes, RNA species and cytoskeleton structures and alters the efficiency of enzymatic reactions in the cell for which the major physiological processes obstacle and creates metabolic imbalance. The physio-morphological description under heat stress reveals reduced rate of germination percentage, plant emergence, abnormal seedlings, poor seedling vigor, reduced radical and plumule growth of germinated seedlings, which are the major impacts caused by heat stress documented in a variety of cultivated plant species. Heat spell at reproductive developmental stages declines flower production or flowers may not produce fruit or seed. In fact, the overproduction of reactive oxygen species (ROS) causing wide-ranging cellular damage and inhibition of physiological functions in plants. Nevertheless, stress positively leads to induce Ca^{2+} influx and cytoskeletal restructuring, ensuing in the upregulation of mitogen activated protein kinases (MAPK) and calcium dependent protein kinase (CDPK). Signaling of these cascades at nuclear level that leads to the production of antioxidants and compatible osmolytes for cell water balance and osmotic adjustment. The sensing of high temperature and induction of signaling cascades are vital adaptive ladder in managing with challengers of heat stress. Significantly, thermotolerance mechanism is largely connected to display of heat shock response and this is completed by reprogramming of gene expression; thereby allowing plants to manage with heat stress. For that reason, greater emphasis on heat stress management is obligatory for heat tolerance features.

Keywords: Heat stress, Thermotolerance, Signaling, Tolerance

INTRODUCTION

Heat stress is a major abiotic stress that influences plant morphology, physiology, reproduction and productivity worldwide. Plant growth and development inhabits repetitive biochemical reactions that are susceptible to temperature. Heat stress, singly or in combination with drought, is a common constraint during anthesis and grain filling stages in many cereal crops of temperate regions. Even a short period of heat stress can cause significant decrease in floral buds and flowers abortion though great variations in sensitivity within and among plant species and variety exist. The escalating drastic impacts of heat stress (HS) are putting global food production as well as nutritional security at high risk. The physiological and molecular responses to HS are vital to research areas. Molecular techniques are being

adopted for producing heat tolerant crop plants. With a view to successfully produce heat stress tolerant crop varieties under the condition of global climate change, there is acute need of knowledge and investigations about heat stress

Corresponding author: Hemantaranjan A, Department of Plant Physiology, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi - 221005, India, Tel: 091 9415302653; E-mail: hemantaranjan@gmail.com

Citation: Hemantaranjan A, Bhanu AN & Lalotra S. (2021) Plant Responses to Heat Stress and Thermotolerance. J Agric Forest Meteorol Res, 4(2): 351-356.

Copyright: ©2021 Hemantaranjan A, Bhanu AN & Lalotra S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

tolerance mechanisms at physiological, biochemical and molecular levels. Plants continuously struggle to modify their metabolic process in many ways in response to heat stress, specifically by generating key solutes that leads to establish proteins and osmotic adjustment and re-establish the redox balance of cell and homeostasis by modifying the antioxidant system. There is an immense need to systematically assess wild species and accessions tolerating extreme degrees of higher temperatures. In that regard, searching for novel donors with high heat tolerance or escape mechanisms is of key importance.

Increased food demand with crops growing under sub-optimal conditions. Consequently, it is genuine to mention that during crop growth and development, temperature plays an important role in dry matter partitioning, transpiration [1,2] photosynthetic activity, respiration [3,4] and root and plant development [5].

In this article, the impacts of heat stress and tolerance at certain growth and developmental levels of plants and approaches that are used to improve heat stress tolerance in crop plants have been reviewed briefly, whereas detailing molecular advances made in heat stress research in the last several years are beyond the scope of this review article.

HEAT STRESS RESPONSES

High temperature stress is a major environmental stress that influences physiology of plants and limits plant growth, metabolism and productivity worldwide. Heat stress differentially influences the constancy of various proteins, membranes, RNA species and cytoskeleton structures and alters the efficiency of enzymatic reactions in the cell for which the major physiological processes obstacle and creates metabolic imbalance [6-9]. The ideal conditions for plant growth and development generally occur within a different range of temperature [10] with low or high temperature (HT) reducing growth and developmental rates [11,12]. Similarly, heat stress has far-reaching effects on plant reproductive organ, seed weight and number of seeds. The increase in temperatures to crop-specific thresholds (for wheat 10-21°C) enhance the rate of grain filling by enhancing cell-division rates in the tissue of endosperm and increasing rates of metabolism [13]. Dry matter partitioning, which is the product of the movement of photosynthetic assimilate from source-sink organs, is enhanced between 10-30°C in winter-season cereals [14]. The devastating decrease in growth and development, harvest index (HI) and seed yield were found for various crops. The decline of grain number resulted from the impact of heat stress on meiosis followed by transfer of pollen during anthesis; growth of ovaries during pre-anthesis periods.

The regulation of heat-shock responses in inflorescence is largely uncharacterized [15,16] as male and female organs are most sensitive to extreme temperature, especially $\geq 30^{\circ}\text{C}$ [17]. Besides these, heat stress damages both male and

female gametophytes and as a consequence, there is decreased pollen viability, reduced pollen germination, pollen tube growth inhibition, stigma receptivity reduction and reduced ovule function, declined fertilization, limited embryogenesis, poor ovule viability, enhanced ovule abortion and a decrease in yield.

PHYSIOLOGICAL RESPONSES

The physio-morphological characteristics under heat stress reduced germination percentage, plant emergence, abnormal seedlings, poor seedling vigor, reduced radical and plumule growth of germinated seedlings are major impacts caused by heat stress recognized in a variety of cultivated plant species [18-20]. Damage to leaf-tip and margins, rolling and drying of leaves and necrosis were observed in sugarcane due to heat stress [21]. Major alterations occur in chloroplasts like altered structural organization of thylakoids, loss of grana stacking and swelling of grana under heat stress [22,23]. Further, all through reproduction, a short period of heat stress can cause significant decrease in floral buds and flowers abortion although great variations in sensitivity within and among plant species and variety exists [24]. Even heat spell at reproductive developmental stages plant may produces no flowers or flowers may not produce fruit or seed [25,26].

Membrane damage

Heat stress severely affects the structure and functions of the membrane, thereby increasing membranes fluidity due to denaturation of proteins and increased level of unsaturated fatty acids, causing a transition from solid gel to flexible crystalline liquid structure [27]. An electrolyte leakage value acts as a pointer of membrane injury that reflects stress-induced alterations. This has been used to evaluate the thermostability of membranes under heat stress [28]. The increased permeability and leakage of ions out of the cell has been used as a measure of cell membrane stability and as a screen test for heat stress tolerance [29]. This happens due to overproduction of reactive oxygen species (ROS) causing extensive cellular damage and inhibition of physiological processes in plants. Although anti-oxidative mechanisms would be an immediate endogenic choice of the plants to counter ROS production, this mechanism can be impaired by heat stress causing a rise in ROS intracellular concentration and an increase in the damage. Diverse metabolic pathways are reliant upon enzymes, which are responsive to various degrees of high temperatures. The accumulation of surplus and damaging ROS are most commonly singlet oxygen ($^1\text{O}_2$), superoxide radical (O_2^-), hydrogen peroxide (H_2O_2) and hydroxyl radical ($\text{OH}^\cdot$) which are responsible for oxidative stress [30]. The major sites of ROS generation are the reaction centers of PSI and PSII in chloroplasts though ROS are also generated in other organelles, viz., peroxisomes and mitochondria [31]. Researchers identified tolerant genotypes that are proved to be more productive under extreme field stress conditions. The thermostability of

the membrane has been successfully employed to evaluate HS tolerance in several crops worldwide.

Photosynthesis

Photosynthesis is a vital process and is often restricted by various abiotic stresses especially under the heat stress and high light conditions. The negative impact of heat stress on plant growth and crop yield were mainly caused by its negative impacts on the photosynthetic process, which are the most thermosensitive aspects of plant functions [32]. The relative water content (RWC), chlorophyll content and PSII activity decreased under high light and heat co-stresses [33]. It is found that photosystem II is thought to be more highly responsive to HS or high light than photosystem I [34]. The photosynthetic process is very sensitive under HS conditions and reduction in chlorophyll contents might be one of the main reasons for the decline in photosynthesis, as an enzyme chlorophyllase helps in conversion of chlorophyll into phytol and chlorophyllide [35]. Remarkably, the increase in temperature from 18-33°C raises the rates of maintenance respiration by greater than 80% [36]. Under elevated temperatures, the rate of respiration measurement could be an appropriate pointer for stimulation of plant response to heat stress, as the rate of respiration rises much more than the rate of photosynthesis initially decreases [10].

Water relations

Heat stress influences plant-water relations due to the faster depletion of water from the soil. This largely affects the temperature of soil and the overall transpiration [37]. Heat stress evidently directly and indirectly affects plant functions and leads to osmotic adjustments by impaired photosynthesis, enhanced respiration, a decline in leaf osmotic potentials and decreased sugar concentration level [38]. Water loss during daytime was more common under heat stress, as of increased transpiration than night time, causing stress in snap bean (*Phaseolus vulgaris* L.) [39]. Under severe heat stress, high stomatal conductance boosts transpirational heat dissipation intolerant genotypes of chickpea as long as soil water is available [40].

PLANT TOLERANCE TO HEAT STRESS

High temperature effects on plant growth and development are reliant upon plant species. With an increasing climate change scenario, there is a better possibility of air temperatures exceeding the optimum range for many species. Exposure of plants to temperature extremes at the onset of the reproductive stage has a major impact on fruit or grain production across all species. Heat stress disturbs cellular homeostasis and causes denaturation and dysfunction in many proteins, leading to severe retardation in growth, development and even death. In plants, the major sites of heat stress injury are the oxygen-evolving complex (OEC) along with associated biochemical reactions in photosystem II (PSII). Ultimately, efficiency of electron transport is reduced or altered affecting electron flow from

OEC towards the acceptor side of PSII. These alterations affect the generation of ATP and the regeneration of Rubisco for carbon fixation [41].

AVOIDANCE AND TOLERANCE MECHANISM

Plants show various survival mechanisms under heat stress, which include long-term morphological and phenological adaptations and short-term acclimation or avoidance mechanisms such as transpirational cooling, alteration of leaf orientation or changing of membrane lipid compositions. Stomatal closure and water loss reduction, enhanced trichomatous and stomatal densities and bigger xylem vessels are the main heat-induced features in crop plants [42]. The capability of the plant to develop and grow to produce economic yield under heat stress is known as thermotolerance. Some important mechanisms including ion transporter, late embryogenesis abundant proteins (LEA), antioxidant defense, osmoprotectants and some factors linked with signaling cascade and transcriptional controls are fundamentally important to respond to heat stress [43]. It has been observed that several tissues in crop plants show differences in terms of developmental exposure, complexity and responses toward the prevailing stress types [44]. These mechanisms help to regenerate homeostasis and to protect and repair damaged membranes and proteins [45].

The osmotic or ionic effects, or changes in temperature or membrane fluidity are the initial stress signals, which activate stress-responsive mechanisms to re-establish homeostasis and protect and repair damaged proteins and membranes. Heat stress accelerates the kinetic energy and movement of molecules across membranes thereby loosening chemical bonds within molecules of biological membranes. This makes the lipid bilayer of biological membranes more fluid by either denaturation of proteins or an increase in unsaturated fatty acids [46]. The tertiary and quaternary structures of membrane proteins are also distorted by heat stress. Such distortions develop the permeability of membranes negatively and as a result there is increased loss of electrolytes. The enhanced solute leakage is an indication of decreased cell membrane thermostability (CMT). This indirect measure of heat-stress tolerance in diverse plant species of electrolyte leakage is influenced by plant/tissue age, sampling organ, developmental stage, growing season, amount of hardening and plant species. Heat tolerance in some species does not correlate with the degree of lipid saturation.

The more fluidity of lipid bilayer under stress leads to the initiation of Ca^{2+} influx and cytoskeletal restructuring, consequential in the upregulation of mitogen activated protein kinases (MAPK) and calcium dependent protein kinase (CDPK). Signaling of cascades at nuclear level shows the way to the production of antioxidants and compatible osmolytes for cell water balance and osmotic adjustment. Further, the production of ROS in the organelles (e.g. chloroplast and mitochondria) is of great significance for

signaling as well as production of antioxidants [47]. The antioxidant defense mechanism is a part of heat-stress adaptation and its strength is correlated with acquisition of thermotolerance [48]. In short, sensing of high temperature and induction of signaling cascades are important adaptive steps in coping with challenges of heat stress.

HEAT SHOCK PROTEINS

Thermotolerance mechanism is mainly related to display of heat shock response and completed by reprogramming of gene expression and thereby allowing plants to cope with the heat stress. During heat stress, the mRNAs encoding non-heat stress induced proteins are destabilized. Using *Arabidopsis* mutants, it was shown that, apart from heat shock proteins (HSP32 and HSP101), ABA, ROS and SA pathways are involved in the development and maintenance of acquired thermotolerance [49].

Under heat stress, most of the stress proteins are soluble in water and consequently contribute to stress tolerance most probably via hydration of cellular structures [50,51] elucidated the fact that during heat stress HSPs, ranging in molecular mass from about 10 to 200 kDa, have chaperone like functions and are involved in signal transduction. The tolerance conferred by HSPs results in improved physiological phenomena such as photosynthesis, assimilate partitioning, water and nutrient use efficiency and membrane stability [52-54]. Interestingly, in certain crop genotypes or species, such improvements make plant growth and development possible under heat stress.

CONCLUSION

In view of foreseen global warming, greater emphasis on heat stress management is obligatory for heat tolerance features. In addition, knowledge relating to molecular basis and mechanism of tolerance that includes molecular cloning and characterization of genes is considered to pave the way for engineering plants that can withstand heat stress.

REFERENCES

1. Crawford AJ, McLachlan DH, Hetherington AM, Franklin KA (2012) High temperature exposure increases plant cooling capacity. *Curr Biol* 22: 396-397.
2. Zhao J, Hartmann H, Trumbore S, Ziegler W, Zhang Y (2013) High temperature causes negative whole-plant carbon balance under mild drought. *New Phytol* 200: 330-339.
3. Sage RF, Zhu XG (2011) Exploiting the engine of C4 photosynthesis. *J Exp Bot* 62: 2989-3000.
4. Atkin OK, Tjoelker MG (2003) Thermal acclimation and the dynamic response of plant respiration to temperature. *Trends Plant Sci* 8: 343-351.
5. Wolkovich EM, Cook BI, Allen JM, Crimmins TM, Betancourt JL, et al. (2012) Warming experiments under predict plant phenological responses to climate change. *Nature* 485: 494-497.
6. Ruelland E, Zachowski A (2010) How plants sense temperature. *Environ Exp Bot* 69: 225-232.
7. Suzuki N, Miller G, Morales J, Shulaev V, Torres MA, et al. (2011) Respiratory burst oxidases: The engines of ROS signaling. *Curr Opin Plant Biol* 14: 691-699.
8. Suzuki N, Koussevitzky S, Mittler R, Miller G (2012) ROS and redox signaling in the response of plants to abiotic stress. *Plant Cell Environ* 35: 259-270.
9. Pagamas P, Nawata E (2008) Sensitive stages of fruit and seed development of chilli pepper (*Capsicum annuum* L var. Shishito) exposed to high-temperature stress. *Sci Hort* 117: 21-25.
10. Criddle RS, Smith BN, Hansen LD (1997) A respiration based description of plant growth rate responses to temperature. *Planta* 201: 441-445.
11. Sánchez B, Rasmussen A, Porter JR (2014) Temperatures and the growth and development of maize and rice: A review. *Glob Chang Biol* 20: 408-417.
12. Ciais P, Reichstein M, Viovy N, Granier A, Ogée J, et al. (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437: 529-533.
13. Barnabas B, Jager K, Feher A (2008) The effect of drought and heat stress on reproductive processes in cereals. *Plant Cell Environ* 31: 11-38.
14. Marcelis LFM (1996) Sink strength as a determinant of dry matter partitioning in the whole plant. *J Exp Bot* 47: 1281-1291.
15. Orbovic V, Poff KL (2007) Effect of temperature on growth and phototropism of *Arabidopsis thaliana* seedlings. *J Plant Growth Regul* 26: 222-228.
16. Van Der Ploeg A, Heuvelink E (2005) Influence of sub-optimal temperature on tomato growth and yield: A review. *J Hort Sci Biotechnol* 80: 652-659.
17. Yang LY, Yang SL, Li JY, Ma JH, Pang T, et al. (2018) Effects of different growth temperatures on growth, development and plastid pigments metabolism of tobacco (*Nicotiana tabacum* L.) plants. *Bot Stud* 59: 5.
18. Kumar S, Kaur R, Kaur N, Bhandhari K, Kaushal N, et al. (2011) Heat-stress induced inhibition in growth and chlorosis in mung bean (*Phaseolus aureus* Roxb) is partly mitigated by ascorbic acid application and is related to reduction in oxidative stress. *Acta Physiol Plant* 33: 2091-2101.
19. Piramila BHM, Prabha AL, Nandagopalan V, Stanley AL (2012) Effect of heat treatment on germination seedling growth and some biochemical parameters of

- dry seeds of black gram. Int J Pharm Phytopharmacol Res 1: 194-202.
20. Toh S, Imamura A, Watanabe A, Okamoto M, Jikumaru Y, et al. (2008) High temperature-induced abscisic acid biosynthesis and its role in the inhibition of gibberellin action in Arabidopsis seeds. Plant Physiol 146: 1368-1385.
 21. Omae H, Kumar A, Shono M (2012) Adaptation to high temperature and water deficit in the common bean (*Phaseolus vulgaris* L.) during the reproductive period. J Bot.
 22. Rodríguez M, Canales E, Borrás-Hidalgo O (2005) Molecular aspects of abiotic stress in plants. Biotechnol Appl 22: 1-10.
 23. Ashraf M, Hafeez M (2004) Thermotolerance of pearl millet and maize at early growth stages: Growth and nutrient relations. Biol Plant 48: 81-86.
 24. Sato S, Kamiyama M, Iwata T, Makita N, Furukawa H, et al. (2006) Moderate increase of mean daily temperature adversely affects fruit set of *Lycopersicon esculentum* by disrupting specific physiological processes in male reproductive development. Ann Bot 97: 731-738.
 25. Maheswari M, Yadav SK, Shanker AK, Kumar MA, Venkateswarlu B (2012) Overview of plant stresses: Mechanisms adaptations and research pursuit. In: Venkateswarlu B, Shanker AK, Shanker C, Maheswari M, editors, Crop Stress and Its Management: Perspectives and Strategies. Springer Dordrecht: The Netherlands, pp: 1-18.
 26. Foolad MR (2005) Breeding for abiotic stress tolerances in tomato. In: Ashraf M, Harris PJC, editors, Abiotic Stresses: Plant Resistance through Breeding and Molecular Approaches. The Haworth Press Inc.: New York, NY, USA, pp: 613-684.
 27. Wahid A, Gelani S, Ashraf M, Foolad MR (2007) Heat tolerance in plants: An overview. Environ Exp Bot 61: 199-223.
 28. Xu S, Li J, Zhang X, Wei H, Cui L (2006) Effects of heat acclimation pre-treatment on changes of membrane lipid peroxidation, antioxidant metabolites and ultrastructure of chloroplasts in two cool-season turf grass species under heat stress. Environ Exp Bot 56: 274-285.
 29. El Basyoni I, Saadalla M, Baenziger S, Bockelman H, Morsy S (2017) Cell membrane stability and association mapping for drought and heat tolerance in a worldwide wheat collection. Sustainability 9: 1606.
 30. Asada K (2006) Production and scavenging of reactive oxygen species in chloroplasts and their functions. Plant Physiol 141: 391-396.
 31. Soliman WS, Fujimori M, Tase K, Sugiyama SI (2011) Oxidative stress and physiological damage under prolonged heat stress in C3 grass *Lolium perenne*. Grassland Sci 57: 101-106.
 32. Wang D, Heckathorn SA, Mainali K, Tripathee R (2016) Timing effects of heat-stress on plant ecophysiological characteristics and growth. Front Plant Sci 7: 1-11.
 33. Chen Y, Zhang Z, Tao F, Palosuo T, Rötter RP (2017) Impacts of heat stress on leaf area index and growth duration of winter wheat in the North China Plain. Field Crop Res, pp: 230-237.
 34. Zivcak M, Brestic M, Kalaji HM, Govindjee (2014) Photosynthetic responses of sun- and shade-grown barley leaves to high light: Is the lower PSII connectivity in shade leaves associated with protection against excess of light? Photosynth Res 119: 339-354.
 35. Mishra D, Shekhar S, Agrawal L, Chakraborty S, Chakraborty N (2017) Cultivar-specific high temperature stress responses in bread wheat (*Triticum aestivum* L.) associated with physicochemical traits and defence pathways. Food Chem 221: 1077-1087.
 36. Penning De Vries FWT (1975) The cost of maintenance processes in plant cells. Ann Bot 39: 77-92.
 37. Sita K, Sehgal A, Hanumantha Rao B, Nair RM, Vara Prasad PV, et al. (2017) Food legumes and rising temperatures: Effects, adaptive functional mechanisms specific to reproductive growth stage and strategies to improve heat tolerance. Front Plant Sci 8: 1-30.
 38. Prasad PVV, Staggenborg SA, Ristic Z, Ahuja LR, Reddy VR (2008) Impacts of drought and/or heat stress on physiological, developmental, growth and yield processes of crop plants. In: Response of Crops to Limited Water: Understanding and Modeling Water Stress Effects on Plant Growth Processes. American Society of Agronomy; Crop Science Society of America; Soil Science Society of America: Madison, WI, USA.
 39. Tsukaguchi T, Kawamitsu Y, Takeda H, Suzuki K, Egawa Y (2003) Water status of flower buds and leaves as affected by high temperature in heat-tolerant and heat-sensitive cultivars of snap bean (*Phaseolus vulgaris* L.). Plant Prod Sci 6: 24-27.
 40. Kaushal N, Awasthi R, Gupta K, Gaur P, Siddique KHM (2013) Heat-stress-induced reproductive failures in chickpea (*Cicer arietinum*) is associated with impaired sucrose metabolism in leaves and anthers. Funct Plant Biol 40: 1334-1349.

41. Tack J, Lingenfelser J, Jagadish SVK (2017) Disaggregating sorghum yield reductions under warming scenarios exposes narrow genetic diversity in US breeding programs. *Proc Natl Acad Sci U S A* 114: 9296-9301.
42. Srivastava S, Pathak AD, Gupta PS, Kumar A, Kumar A (2012) Hydrogen peroxide-scavenging enzymes impart tolerance to high temperature induced oxidative stress in sugarcane. *J Environ Biol* 33: 657-661.
43. Sehgal A, Sita K, Nayyar H (2016) Heat stress in plants: Sensing and defence mechanisms. *J Plant Sci Res* 32: 195-210.
44. Queitsch C (2000) Heat shock protein 101 plays a crucial role in thermotolerance in Arabidopsis. *Plant Cell* 12: 479-492.
45. Vinocur B, Altman A (2005) Recent advances in engineering plant tolerance to abiotic stress: Achievements and limitations. *Curr Opin Biotechnol* 16: 123-132.
46. Savchenko GE, Klyuchareva EA, Abrabchik LM, Serdyuchenko EV (2002) Effect of periodic heat shock on the membrane system of etioplasts. *Russ J Plant Physiol* 49: 349-359.
47. Bohnert HJ, Gong Q, Li P, Ma S (2006) Unravelling abiotic stress tolerance mechanisms—getting genomics going. *Curr Opin Plant Biol* 9: 180-188.
48. Maestri E, Klueva N, Perrotta C, Gulli M, Nguyen HT, et al. (2002) Molecular genetics of heat tolerance and heat shock proteins in cereals. *Plant Mol Biol* 48: 667-681.
49. Larkindale J, Huang B (2005) Effects of abscisic acid salicylic acid ethylene and hydrogen peroxide in thermotolerance and recovery for creeping bent grass. *Plant Growth Regul* 47: 17-28.
50. Wahid A (2007) Physiological implications of metabolites biosynthesis in net assimilation and heat stress tolerance of sugarcane (*Saccharum officinarum*) sprouts. *J Plant Res* 120: 219-228.
51. Schoffl F, Prandl R, Reindl A (1999) Molecular responses to heat stress. In: Shinozaki K, Yamaguchi-Shinozaki K, Eds, *Molecular Responses to Cold, Drought, Heat and Salt Stress in Higher Plants*. RG Landes Co.: Austin, Texas, pp: 81-98.
52. Camejo D, Rodriguez P, Morales MA, Dellamico JM, Torrecillas A (2005) High temperature effects on photosynthetic activity of two tomato cultivars with different heat susceptibility. *J Plant Physiol* 162: 281-289.
53. Ahn YJ, Zimmerman JL (2006) Introduction of the carrot HSP177 into potato (*Solanum tuberosum* L) enhances cellular membrane stability and tuberization *in vitro*. *Plant Cell Environ* 29: 95-104.
54. Momcilovic I, Ristic Z (2007) Expression of chloroplast protein synthesis elongation factor EF-Tu in two lines of maize with contrasting tolerance to heat stress during early stages of plant development. *J Plant Physiol* 164: 90-99.