Iournal of Clinical Trials & Research

JCTR, 1(2): 17-20 www.scitcentral.com



Original Review: Open Access

Why Reconstructive Neurosurgery?

Visocchi M*

*Institute of Neurosurgery Catholic University of Rome, Italy

Received April 11, 2018; Accepted April 29, 2018; Published October 20, 2018

ABSTRACT

Spine and peripheral nerves reconstructive surgery, central nervous system revascularization (surgical, radiointerventional), neuromodulation, bioengineering and transplantation are the recent tools to promote reconstruction, restoration and rehabilitation.

The ISRN is an "open" multidisciplinary Society in evolution. Many different souls inspire new trends in many different neurosurgical fields, all dealing with "neurosurgical reconstruction". Spinal, functional, vascular, radio and oncologic neurosurgeons are the addressees of our proposal along with biologists, bioengineers, anatomists physiologists, physiotherapists who are precious and irreplaceable inspirers. These are the three key words of our creed and all fulfill the aim of ISRN, dealing with mechanical morphological and functional restoration.

Keywords: Neurosurgical reconstruction, Neurosurgical restoration, Neurosurgical rehabilitation

INTRODUCTION

Founded in 1955, the WFNS is a professional, scientific, non-governmental organization comprising 5 Continental Associations, 115 National Neurosurgical Societies and 7 Affiliate Societies, representing some 30,000 neurosurgeons worldwide. The World Federation of Neurosurgical Societies (WFNS) aspires to promote global improvement in neurosurgical care, training and research to benefit the patients. The WFNS is governed by an Executive Committee consisting of two Delegates from each Member Society and an Administrative Council composed of the Officers of the Federation, who are elected every four years. The EC meets every two years and is guided by the Administrative Council which meets at least annually.

The goals of the WFNS are deliberated and pursued through scientific, standing and ad hoc committees and during the International Congress of Neurological Surgery which takes place every four years. The Neurorehabilitation and Reconstructive Neurosurgery Committee (NRNC) is a special Section which promotes all those activities of the WFNS aiming at implement and promote all the restorative, reconstructive and augmentative neurosurgical procedures, grossly identified in the past as the Functional Neurosurgery subspecialty but now updating, evolving and merging with neuro oncology, spine surgery neuroradiological and neurophysiological intraoperative assistance and new technologies lato sensu. As matter of fact the Neuromodulation Committee is a different Section and it is administered by a completely different Board working in an independent way and pursuing different objectives. Two years after the institution of the WFNS Committee, the members of the Board of the Neurorehabilitation and Reconstructive Neurosurgery Committee felt the need to found a new Society, more free to deal with other Medical and /or Surgical Societies and expert researchers of different branches of biology, physiology, physiotherapy. The IV International Congress of International Society of Reconstructive Neurosurgery (ISRN) along with the VII Neurorehabilitation and Reconstructive Neurosurgery Symposium (WFNS) was held in Cerveteri (Rome) on SEPTEMBER the 12th - 14th, 2015. The President of the Congress was Prof Franco TOMASELLO, former Rector of the University of Messina Italy and actually Vice President of WFNS and myself Massimiliano VISOCCHI, Past President of the Italian Society of Neurosonology and Cerebral Haemodinamics, former Secretary of the Spine Section of the Italian Society of Neurosurgery, Associate Professor in Neurosurgery at the Catholic University of

*Corresponding Author: Visocchi M, Institute of Neurosurgery Catholic +39-360807781; Email: University of Rome, Tel: Italy, mvisocchi@hotmail.com

Citation: Visocchi M. (2018) Why Reconstructive Neurosurgery? J Clin Trials Res, 1(2): 17-20.

Copyright: ©2018 Visocchi M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

17 J Clin Trials Res (JCTR)

J Clin Trials Res, 1(2): 17-20 Visocchi M

of Rome and Visiting Professor at the Shanghai Jiao Tong University School of Medicine. In such an occasion I was appointed President Elect of the ISRN and soon after I started to ask to myself some more questions dealing with Reconstructive Neurosurgery and also to try to better investigate on the true roots of our Society and the supposed mission of ISRN. In other words "were we come from? Who are we and where are we going?"

Now, also as Chairman Elect of the NRNC my main impulse is to have a look on the current literature on the topic. If we search for "Reconstructive Neurosurgery" on Pub Med. Gov the US National Library of Medicine National Institute of Health we can find 3849 papers updated at April 2018 the 11th. The papers are spanning from the latter, printed on April 2018 and harbouring a very intriguing title: Independent factors affecting postoperative complication rates after custom-made porous hydroxyapatite cranioplasty: a single-center review of 109 cases [1] to the former, the very first recognized paper, printed on November 1947 and dealing with A report of the early results in tantalum cranioplasty [2]. In between these papers there are many others dealing with craniofacial endoscopy [3], spinal instrumentation and fusion procedures [4], perpheral nerves reconstructive procedures [5], vascular reconstructive and cerebral blood flow restorative surgical procedures [5], neuronavigation and video assisted neurosurgical procedures [6], craniofacial complex surgical procedures [7], intraoperative neuromonitoring Surprisingly many paper dealing with genetics, molecular and cytochemical studies dealing with central and peripheral nervous system are available as well [9]. Reconstruction is intended in a pure mechanical way and no concepts dealing with restoration or rehabilitation arise from the lines of the majority of the manuscripts but just a simple exposition of surgical techniques and procedures aiming at just repairing something. More in details even "dysfunctional" syndromes as Raynaud syndrome are faced with mechanical procedures instead of neuroaugmentative ones [10].

Neurosurgeons know very well that central nervous system, along with the spine and the skull bone components, has an intrinsic pattern of complex physiological nature both from the neuro - chemical and the biomechanical point of view. Starting from the Neurorehabilitation Committee of the WFNS experience, we turned our gaze toward restoration and rehabilitation very soon and now, since ISRN is an "open" multidisciplinary Society in evolution, we first intended neurosurgical reconstruction, latu sensu, in a new way: reconstruction means also rehabilitation and restoration.

Nevertheless in the common literature restoration is strongly linked to the concept of reconstruction as demonstrated by the title of the following paper included in the Pub Med list of Reconstruction: Restoration of the orbital aesthetic subunit with the thoracodorsal artery system of flaps in patients undergoing radiation therapy [11].

Otherwise neurorehabilitation has different sounds and significances in the culture of the neurosurgeons; although it still deal with reconstruction e. g Functional restoration of diaphragmatic paralysis: an evaluation of phrenic nerve reconstruction [12] and also with replantation as descripted in in the paper Six years of follow-up after bilateral hand replantation [13], functional neurorehabilitation merges with the concept of neuromodulation and neuromodulation is the core of the functional neurosurgery. So more simply spine and peripheral nerves reconstructive surgery, central nervous system revascularization (surgical, radiointerventional), neuromodulation, bioengineering and transplantation are the recent tools to promote reconstruction in the special sense intended by our Society.

More in details neuromodulation classically deals with is the physiological process by which a given neuron uses one or more neurotransmitters to regulate diverse populations of neurons. This is in contrast to classical synaptic transmission, in which one presynaptic neuron directly influences a single postsynaptic partner. Neuromodulators secreted by a small group of neurons diffuse through large areas of the nervous system, affecting multiple neurons. Otherwise in the surgical praxis the meaning shifted toward all the surgical tools armamentarium dealing with all the procedures involved in CNS electrical and chemical stimulation as performed with spinal cord stimulation (SCS) deep brain stimulation (DBS), cortical brain stimulation (CBS) drug delivery systems implantation (DDS) aimed at facing with pain, movement disorders, spasticity, bowel and bladder dysfunction, peripheral heart and cerebral vasculopaties. Interestingly an interpherence of SCS was first reported by Hosobuchi in 1986 also with CBF; he reported the intriguing effect of SCS on CBF in human beings, along with the demonstration that spinal cord stimulation can improve peripheral blood flow. Following these initial clinical and experimental observations, he first described the use of cervical SCS for the treatment of cerebral ischemia in man in 1991 [14].

Others have reported that SCS improves clinical symptoms of patients in persistent vegetative states, improves CBF in stroke patients, suppresses the hemodynamic mechanism underlying headache attacks in migraneous patients, and increases locoregional blood flow in high grade brain tumours in humans. In animals, SCS has been shown to prevent progression of cerebral infarction, reduce infarct volume, reduce ischemic brain edema and improve vasospasm [15]. Studies of our group have produced variable results: SCS can produce an increase of CBF, a reduction or no effect. In patients studied with both single photon emission cerebral tomography (SPECT) flowmetry

J Clin Trials Res, 1(2): 17-20 Visocchi M

and Transcranial Doppler Sonography (TCD) the size of the induced variations, when present in both, was the same. Cervical stimulation more frequently produces an increase in CBF (61% of cervical stimulations) [16-18]. Experimental studies of our group confirmed that SCS 1) interacts with CO2 with the mechanism of regulation of CBF in a competitive way and produce a reversible functional sympathectomy; 2) produces similar flowmetric changes in the brain as well as in the eyes; 3) can improve both clinical and hemodynamic ischemic stroke in humans, 4) prevents hemodynamic deterioration in the experimental combined ischemic and traumatic brain injury; and 5) prevents experimental early vasospasm [19-21]. On the other hand, trigeminal ganglion stimulation can have opposite effects [22].

But when we speak about reconstruction we cannot forget spinal cord and spine surgery, both surgical challenges, from a bio functional point of view the former and from a bio mechanical point of view the latter. Instrumentation and fusion procedures from the upper to the lower levels of the spine have been widely published so far [23-28]. Nevertheless the choice of altenative miminimally invasive video assisted surgical routes for spine reconstruction have got also a big deal of interest and opened new perspectives in reconstruction, rehabilitation and restoration which are the three key words of our creed and all fulfill the aim of ISRN, dealing with mechanical morphological and functional restoration [29-35].

Spinal, functional, vascular, radio and oncologic neurosurgeons are the addressees of our proposal along with biologists, bioengineers, anatomists physiologists, physiotherapists who are precious and irreplaceable inspirers.

REFERENCES

- 1. Still M, Kane A, Roux A, Zanello M, Dezamis E, et al. (2018) Independent factors affecting postoperative complication rates after custom-made porous hydroxyapatite cranioplasty: a single-center review of 109 cases. World Neurosurg 18: S1878-8750.
- Lane S, Webster JE (1947) A report of the early results in tantalum cranioplasty. J Neurosurg 4: 526-529.
- Sreenath SB, Recinos PF, McClurg SW, Thorp BD, McKinney KA, et al. (2015) The Endoscopic Endonasal Approach to the Hypoglossal Canal: The Role of the Eustachian Tube as a Landmark for Dissection. JAMA Otolaryngol Head Neck Surg 17: 1-8.
- 4. Lim SH, Jo DJ, Kim SM, Lim YJ (2015) Reconstructive surgery using dual U-shaped rod

- instrumentation after posterior en bloc sacral hemiesection for metastatic tumor: case report. J Neurosurg Spine 31: 1-5.
- Hendry JM, Alvarez-Veronesi MC, Snyder-Warwick A, Gordon T, Borschel GH (2015) Side-To-Side Nerve Bridges Support Donor Axon Regeneration Into Chronically Denervated Nerves and Are Associated With Characteristic Changes in Schwann Cell Phenotype. Neurosurg.
- 6. Takanari K, Araki Y, Okamoto S, Sato H, Yagi S, et al. (2015) Operative wound-related complications after cranial revascularization surgeries. J Neurosurg 3: 1-6.
- Andrews BT, Thurston TE, Tanna N, Broer PN, Levine JP, et al. (2015) A Multicenter Experience With Image-Guided Surgical Navigation: Broadening Clinical Indications in Complex Craniomaxillofacial Surgery. J Craniofac Surg 26: 1136-1139.
- 8. Biscevic M, Biscevic S, Ljuca F, Smrke BU, Ozturk C, Tiric-Campara M (2014) Motor evoked potentials in 43 high risk spine deformities. Med Arch 68: 345-349.
- 9. Lopez-Bertoni H, Lal B, Li A, Caplan M, Guerrero-Cázares H, et al. (2015) DNMT-dependent suppression of microRNA regulates the induction of GBM tumor-propagating phenotype by Oct4 and Sox2. Oncogene 34: 3994-4004.
- 10. Merritt WH (2014) Role and rationale for extended periarterial sympathectomy in the management of severe Raynaud syndrome: techniques and results. Hand Clin 31: 101-20.
- Chanowski EJ, Casper KA, Eisbruch A, Heth JA, Marentette LJ, et al. (2013) Restoration of the orbital aesthetic subunit with the thoracodorsal artery system of flaps in patients undergoing radiation therapy. J Neurol Surg B Skull Base 74: 279-285.
- 12. Kaufman MR, Elkwood AI, Colicchio AR, CeCe J, Jarrahy R, et al. (2014) Functional restoration of diaphragmatic paralysis: an evaluation of phrenic nerve reconstruction. Ann Thorac Surg 97: 260-266
- 13. Marques M, Correia-Sá I, Festas MJ, Silva S, Silva AI, et al. (2013) Six years of follow-up after bilateral hand replantation Chir Main 32: 226-234.
- 14. Hosobuchi Y (1985) Electrical stimulation of the cervical spinal cord increases cerebral blood flow in humans. Appl Neurophisyol 48: 372-376.
- Visocchi M (2008) Neuromodulation of cerebral blood flow by spinal cord electrical stimulation: the role of the Italian school and state of art. J Neurosurg Sci 52: 41-47.
- 16. Visocchi M (2006) Spinal cord stimulation and cerebral haemodynamics. Acta Neurochir Suppl 99: 111-116.

J Clin Trials Res, 1(2): 17-20 Visocchi M

17. Meglio M, Cioni B, Visocchi M, Nobili F, Rodriguez G, et al. (1991) Spinal cord stimulation and cerebral haemodynamics. Acta Neurochir 111: 43-48.

- 18. Visocchi M, Argiolas L, Meglio M, Cioni B, Dal Basso P, et al. (2001) Spinal cord stimulation and early experimental cerebral spasm: the functional monitoring and the preventing effect. Acta Neurochir 143: 177- 185.
- Visocchi M, Tartaglione T, Romani R, Meglio M (2001) Spinal cord stimulation prevents the effects of combined ischemic and traumatic brain injury. An MR study. Stereotact Funct Neurosurg 76: 276-281.
- Visocchi M, Cioni B, Pentimalli L, Meglio M (1994) Increase of cerebral blood flow and improvement of brain motor control following spinal cord stimulation in ischemic spastic hemiparesis. Stereotact Funct Neurosurg 62: 103-107.
- 21. Visocchi M (2008) Sympathetic activity does influence cerebral blood flow J Appl Physiol 105: 1369.
- 22. Visocchi M, Chiappini F, Cioni B, Meglio M (1996) Cerebral autoregulation and trigeminal stimulation. A TCD study. Ster Funct Neurosurg 66: 184-192.
- 23. Visocchi M, Di Rocco F, Meglio M (2003) Craniocervical junction instability: instrumentation and fusion with titanium rods and sublaminar wires. Effectiveness and failures in personal experience. Acta Neurochir 145: 265-272.
- 24. Visocchi M, Pietrini D, Tufo T, Fernandez E, Di Rocco C (2009) Pre-operative irreducible C1-C2 dislocations: intra-operative reduction and posterior fixation. The "always posterior strategy". Acta Neurochir 15: 551-560
- Visocchi M, Fernandez EM, Ciampini A, Di Rocco C (2009) Reducible and irreducible os odontoideum treated with posterior wiring, instrumentation and fusion. Past or present? Acta Neuroch 151: 1265-1274.
- 26. Visocchi M (2009) Pre-operative irreducible C1-C2 dislocations: intra-operative reduction and posterior fixation. The "always posterior strategy". Acta Neurochir 151: 551-559.
- 27. Visocchi M, Conforti G, Roselli R, La Rocca G, Spallone A (2015) From less to maximally invasiveness in cervical spine surgery: A "nightmare" case who deserve consideration. Int J Surg Case Rep 9: 85-88.
- 28. Visocchi M, Conforti G, La Rocca G, Roselli R, Spallone A, Barbagallo G (2014) Invasive and mini-invasive lumbar fusions. Does exist a state of art? An enigma in search of solutions. J Neurosurg Sci 58: 113-117.

- 29. Visocchi M, Masferrer R, Sonntag VHK (1998) Dickman: Thoracoscopic approaches to the thoracic spine. Acta Neurochir 140: 737-744.
- 30. Visocchi M, La Rocca G, Della Pepa GM, Stigliano E, Costantini A, et al. (2014) Anterior video-assisted approach to the craniovertebral junction: transnasal or transoral? A cadaver study. Acta Neurochir 156: 285-292.
- 31. Visocchi M, Doglietto F, Della Pepa GM, Esposito G, La Rocca G, et al. (2011) Endoscope-assisted microsurgical transoral approach to the anterior craniovertebral junction compressive pathologies. Eur Spine J 20: 1518-1525
- 32. Visocchi M, et al. (2011) Video-assisted microsurgical transoral approach to the craniovertebral junction: personal experience in childhood. Childs Nerv Syst.
- 33. Visocchi M, Trevisi G, Iacopino DG, Tamburrini G, Caldarelli M, et al. (2014) Odontoid process and clival regeneration with Chiari malformation worsening after transoral decompression: an unexpected and previously unreported cause of "accordion phenomenon". Eur Spine J.
- 34. Visocchi M, Di Martino A, Maugeri R, González Valcárcel I, Grasso V, et al. (2015) Videoassisted anterior surgical approaches to the craniocervical junction: rationale and clinical results. Eur Spine J.
- 35. Visocchi M (2015) Transnasal and transoral approach to the clivus and the craniovertebral junction. J Neurosurg Sci.