

Rich Genetic Diversity of Capsicum Species in Northeast India, as a Potential Source for Chilli Crop Improvement

Karma L Bhutia¹, Nangsol D Bhutia² and VK Khanna^{1*}

¹School of Crop Improvement, College of Post Graduate Studies, CAU (Imphal), Umiam, Meghalaya-793103, India

²Department of Vegetable Sciences, Indian Agricultural Research Institute (IARI), New Delhi-110012, India.

Received April 27, 2018; Accepted August 21, 2018; Published March 26, 2019

ABSTRACT

Northeastern India with its diverse topography ranging from plains of Assam and Tripura to mountainous regions of Sikkim and Arunachal Pradesh possesses great diversity of flora and fauna, including varieties of Capsicum species. Although, Northeastern India is rich in genetic diversity of *C. species*, information about biotic and abiotic stresses and the source of resistance, breeding approaches to improve the chilli landraces of NE India, ethno-pharmacological applications, capsaicin/capsaicinoid content and other useful compounds of many landraces are still largely unexplored. This review article highlights about the various research work carried out so far in chillies of NE India, related to genetic diversity study and tissue culture techniques, etc.

Keywords: Capsicum species, Northeast India, Genetic diversity, Landraces, Capsaicin

Abbreviations: NE India: Northeast India; *C. species*: Capsicum species; CAU: Central Agricultural University

INTRODUCTION

Chillies belong to the genus *Capsicum* of the Solanaceae family. *Capsicum* is derived from the Greek word 'kapsimo', meaning 'to bite' [1]. There are 25-30 species of *Capsicum* of which 5 species, i.e., *C. annum* L., *C. frutescens* mill, *C. chinense*, *C. baccatum* L. and *C. pubescens* have been domesticated and cultivated [2]. *Capsicum* species are diploids with most of them having 24 chromosome ($n=x=12$), but species with genome size of $2n=2x=32$ [3] and $2n=2x=48$ [4] also have been reported. It is a self-pollinated dicot plant. However, occurrence of cross pollination leads to the formation of variants within the species. The primary centre of origin of chilli is thought to be Mexico (American tropics) with secondary centers in Guatemala and Bulgaria [5]. Chilli is believed to have been introduced to India by Portuguese explorers [1] and to NE India by Christian missionaries [6]. The NE India has a rich natural heterogeneity due to its distinctive geographical location which has been identified as one of the twelve "Genetic Epicentres" for the evolution of world flora. In NE India, several *Capsicum* landraces are traditionally grown, which are distinguished based on their morphological appearance (**Figure 1**), flavor and pungency levels. These are referred to by a number of vernacular names, e.g. Naga King chilli (Bhut Jolokia/Bih Jolokia/Naga Jolokia, Lota Bhut, Dhan Jolokia, Krishna Jolokia, Mem Jolokia, Raja

mircha, U-morok), In Sikkim, they are known as Dalle Khursani (round chillies), Thadey Khursani (erect fruit), Thalo Khursani (vegetable type), Jeerey Khursani (thin fruit), Lamchey Khursani (medium size-less pungent) and so on.

Corresponding author: VK Khanna, School of Crop Improvement, College of Post Graduate Studies, CAU (Imphal), Umiam, Meghalaya-793103, India, E-mail: khannavk@rediff.com

Citation: Bhutia KL, Bhutia ND & Khanna VK. (2019) Rich Genetic Diversity of Capsicum Species in Northeast India, as a Potential Source for Chilli Crop Improvement. J Agric Forest Meteorol Res, 2(2): 77-83.

Copyright: ©2019 Bhutia KL, Bhutia ND & Khanna VK. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1. Flowering and fruit morphology of *Capsicum frutescens* (a, b and c), *C. chinense*/King chilli (d, e and f), *C. annum* (g, h and i) and Dalle khursani (*C. annum*) (j and k).

Capsicum is used as a spice in a variety of cuisine all over the world as a basic ingredient. It is also used as a colorant and flavorant to make the foods more attractive and tasty. The archaeological data suggested or proved that the human civilization has been cultivating Capsicum since the past 7000 years to be used in food recipes for hotness and coloration [7]. Capsicum not only gives attractive color and flavor to the foods but also provides vitamin C, vitamin A, vitamin B complex, vitamin E and minerals like Molybdenum, Manganese, Folate, Potassium and Thiamine. Chilli contains seven times more vitamin C than orange. Beta carotenoids, Vitamin C and vitamin A in chillies are antioxidants that destroy free radicals [8]. Pungency (hotness) of the chilli pepper is due to the presence of capsaicin. It is an active component of chili peppers. Capsaicin is the main capsaicinoid in chilli peppers, followed by dihydrocapsaicin, nordihydrocapsaicin, homodihydrocapsaicin and homocapsaicin [9]. Capsaicin present in chilli pepper is used as a counter irritant in Lumbago (pain in the muscles and joints of the lower back), Neuralgia (pain in the distribution of a nerve or nerves), Rheumatic disorders (chronic, often intermittent pain affecting the joints and/or connective tissue) and non-allergic Rhinitis (inflammation of the inner part of the nose that is not caused by an allergy). It has a tonic and carminative action. In combination with Cinchona, it is used for the treatment of intermittent and lethargic affliction,

tympanitis (pain due to inflammation of the middle ear as well as oozing blisters on the eardrum) and paralysis [2]. The plants have also been used as indigenous medicine for asthma, arthritis, dropsy, colic, diarrhea, toothache and muscle cramp [10].

AREA AND PRODUCTION OF CHILLI IN NE STATES OF INDIA

India is the largest producer, consumer and exporter of chillies in the world [11]. In India, per capita consumption is highest for chillies, among the spices produced. It occupies around 30 % of the area shared among major spice crops of the country [12]. Though chillies are grown all over India, NE states contribute 51.72% of its annual production while having only 8% area under chilli cultivation (Spice Statistics, Spice Board, 2004). This region comprising the states of Arunachal Pradesh, Assam, Manipur, Meghalaya, Mizoram, Nagaland, Sikkim and Tripura is located between 22-29.3° N and 89.7-97.8° E. Among the NE states, Assam has the maximum area under chilli cultivation, and it seems Nagaland is at top among the NE states in terms of production (**Table 1**). NE region of India is an excellent place to find the diverse range of chilli genotypes, because there are numerous heterogeneous landraces of chilli available in this region and these landraces serve as a reservoir of genetic variability for chilli breeders [13].

Table 1. Area and Production of chillies (green) in north-eastern states of India in the year 2015-2016.

S. No.	Name of the State	Year: 2015-2016	
		Area (000' ha)	Production (000' t)
01	Assam	21.31	19.12
02	Nagaland	5.40	40.08
03	Mizoram	9.14	9.33
04	Tripura	2.15	19.80
05	Manipur	0.20	1.51
06	Sikkim	0.09	0.32
07	Arunachal Pradesh	1.2*	1.6**
08	Meghalaya	1.8*	1.1**

** indicates the data of the year 2005-2006

Source: Horticulture at a Glance, 2017. Government of India, Ministry of Agriculture and Farmers Welfare, Horticulture Statistics Division

REPORTS ON ASSESSMENT OF GENETIC DIVERSITY IN CAPSICUM SPECIES OF NE INDIA

For strategic germplasm collection, maintenance, conservation and utilization, sound knowledge of diversity in any plant species is an important requirement [14]. Numerous researches have been conducted on genetic diversity of Capsicum using morphological as well as molecular markers. However, there has been very little research on genetic diversity of Capsicum accessions collected from NE India.

Mohammadi and Prasanna [14] assessed genetic diversity using morphological characters of 52 genotypes of Capsicum which were collected from different regions of NE India. Wide variations were observed in various morphological characters studied which suggested the diverse nature of germplasm collected. The study on genetic relationship of nineteen genotypes belonging to *C. annuum*, *C. frutescens*, *C. chinense* and Bhut Jolokia showed that Bhut Jolokia shared more markers from *C. chinense* than with *C. frutescens*. The average genetic similarity of Bhut Jolokia and *C. chinense* was 0.79 which is closer to the mean genetic diversity within *C. chinense* (0.82) than that within *C. frutescens* (0.85). Based on the presence of both *C. frutescens* and *C. chinense* specific markers in Bhut Jolokia, the study concluded it as an interspecific hybrid between the two species [15]. This study also identified some of the species specific markers which will be useful in species identification as well as for marker assisted breeding of chilli crops. Evaluation of genetic diversity of Naga Jolokia with morphological characters revealed a considerable amount of genetic variability with respect to fruit shape and color in this germplasm [16]. Twenty five chilli genotypes which were from NE region of India, evaluated for phytochemical composition and antioxidant potential showed variations for

capsaicin, oleoresin, phenolics, carotenoids and other antioxidants contents [17]. Genetic differences between Bhut Jolokia, *C. frutescens* and *C. chinense* was revealed when six genotypes of Bhut Jolokia with eight known Capsicum species were characterized by using three nuclear genes, namely; ITS 1, ITS 2 and 5.8 S. However, the study failed to discriminate between *C. frutescens* and *C. chinense* [7]. Cluster analysis of genetic diversity of seven Capsicum landraces of *C. annuum*, *C. frutescens* and *C. chinense* using 10 RAPD showed two major clusters where *C. frutescens* and *C. chinense* genotypes grouped in one cluster apart from the *C. annuum* genotypes. The average genetic diversity was more in *C. frutescens* and *C. chinense* genotypes than in *C. annuum* [18]. The genetic diversity analysis of 53 chilli landraces of NE India using pun 1 locus revealed 79 SNPs and 3 indels in pun1 gene and when SSR markers and morphological traits were used to study diversity, they detected 3-9 alleles for each SSR locus with an average of 5.36 alleles/locus. Cluster analysis based on morphological and SSR markers separated the accessions with campanulate fruits type from those with erect fruits type [19]. When SSR and random amplified microsatellite polymorphism (RAMPO) markers were used to evaluate the genetic diversity of germplasm collected from nine countries which included 4 species of Capsicum, improved lines and 5 landraces from NE India, the dendrogram based on both the markers showed two major clusters and was able to separate *C. annuum* genotypes from others. The study revealed the presence of higher genetic diversity in other *Capsicum* spp. than *C. annuum* genotypes [20]. Genetic diversity based on twelve quantitative characters of 30 chilli (*Capsicum annuum* L.) genotypes collected from different regions of India, majority from Arunachal Pradesh, showed considerable variability among the genotypes for the character studied and some genotypes were identified as

better parents for an efficient hybridization programme of chilli [21]. Protein profiling of 30 genotypes of chilli collected from NE India showed considerable variation in banding pattern of total protein which ranged from 7-19 numbers of bands [22]. Five different ecotypes of Bhut Jolokia collected from different locations of NE India showed significant difference in terms of phenolic content combined with ascorbic acid in the fruits [23]. Diversity analysis by estimating the bioactive compounds present in 72 chilli landraces of NE India showed significant difference in total phenolic and flavonoid content. The result also suggested that *Capsicum frutescens* L. is a potent source of natural antioxidants [13]. Significant intra and inter-specific variations for fruit morphological traits, fruiting habits and 65 fruit metabolites were observed in the collected Capsicum germplasm belonging to three Capsicum species, i.e., *C. chinense* (Bhut jolokia, 63 accessions), *C. frutescens* (17 accessions) and *C. annuum* (56 accessions). Phenotypic diversity of 72 landraces of chilli consisting predominantly of *C. frutescens* collected from different regions of NE India using 29 qualitative traits and 14 quantitative traits revealed the vast intra-specific diversity of *C. frutescens* landraces of NE India [24]. SDS-PAGE analysis of 31 chilli genotypes collected from different regions of NE India showed considerable variation in number of protein bands ranging from 8-18 and a total of 70 protein bands as per Rm values were identified by silver staining [25]. Study on about 30 Bhut Jolokia genotypes to determine the distinctiveness of Bhut Jolokia with the germplasm using SSR markers revealed 100% polymorphism in 9 out of 10 markers [26].

OTHER RESEARCH ACTIVITIES ON CHILLIES OF NE INDIA

Although NE India is rich in genetic diversity of chilli crops, there has been very little research conducted in chilli crops belonging to NE India regions. Therefore, there is a high probability of discovering a number of genes/alleles contributing to resistance against abiotic and biotic stresses, some chemicals/compounds that can be used as medicine and other secondary metabolites that are of high value. Below are some of the researches conducted on chillies of NE India for various purposes. Capsaicin and Capsaicinoid have been estimated from different chilli cultivars of NE India [27-30]. Growing environment, sowing time and crop geometry influenced the synthesis of capsaicin or capsaicinoids in Chilli pepper [31,32]. Comparative expression analysis (through qRT-PCR) of candidate genes involved in capsaicin/capsaicinoid biosynthesis pathway showed many fold higher expression of majority of the genes in *C. chinense* compared to *C. frutescens* and *C. annuum* suggesting that the possible reason for extremely high pungency might be due to the higher level of candidate gene(s) expression [33]. Many components of chilli fruits are known to have medicinal values. The ethno-pharmacological applications of king chilli and its derivatives is well described [34-36]. Susceptibility of the

chilli genotypes to a number of abiotic and biotic stresses has restricted their potential yield. Abiotic stresses which significantly reduced the yield and quality of peppers include extreme of temperature, moisture, light, nutrients and pH among others. Biotic factors include susceptibility of chilli plants to various fungi, bacteria, viruses, nematodes and various insects. Among the biotic factors, the one that poses a great threat to cultivated chilli is viral infection [37]. All these problems can be overcome by identifying novel sources of resistance [38-40], heterosis breeding [41] and proper identification and management of causal organism [42-46]. Number of characteristics like very low germination percentage, recalcitrant nature, genotype dependent, etc. associated with the Capsicum creates problems in improvement of chilli plants by conventional breeding techniques as compared to other solanaceous crops [2]. Hence, the plant biotechniques which include plant tissue culture and genetic engineering are gradually becoming a functional aspect of classical breeding programs and boost the improvement in Capsicum crop. Many researchers [4,23,28,32,47-57] reported the efficient method for *in vitro* regeneration/micro-propagation of chillies belonging to NE India, that will help in further improvement of the chilli crop through biotechnological approaches that involve tissue culture.

CONCLUSION

The NE India has a rich natural heterogeneity due to its distinctive geographical location where several landraces of crops including Capsicum species are traditionally grown. These landraces of chillies have various traditional uses including uses in different ethnic cuisines and ethno-pharmacological use [58]. Although many studies have been done in chilli landraces of NE India including diversity/evolution, *in vitro* regeneration, scientific cultivation, diseases and their management, the potentiality of chilli landraces of NE India as the source of biotic and abiotic stress resistance, medicinal properties, etc. are still largely unexplored. Chilli cultivation in NE India also faces the problems like lack of warehouse/godown for proper storage, poor transport facility for harvested product and lack of market information to farmers for efficient and timely production of chilli on their farms. There is a need to conduct more researches on NE India chilli to develop varieties that are better yielding, tolerant to various stresses, etc. Further, awareness about market situation, cultivation techniques, proper storage and transportation will make chilli cultivation more remunerative for chilli farmers in NE India [59-61].

ACKNOWLEDGEMENT

We thank Dean, College of Post-Graduate Studies and Chairman, School of Crop Improvement, CAU (Imphal) for support. Bhutia KL and Bhutia ND are grateful to UGC, Government of India for National Fellowship.

REFERENCES

- Basu SK, De AK (2003) Capsicum: Historical and botanical perspectives. In: De AK (ed.) The genus Capsicum. Taylor & Francis, London, pp 1-15.
- Kothari SL, Joshi A, Kachhawaha S, Ochoa-Alejo N (2010) Chilli pepper - A review on tissue culture and transgenics. Biotechnol Adv 28: 35-48.
- Wang D, Bosland PW (2006) The genes of Capsicum. Horticult Sci 41: 1169-1187.
- Dafadar A, Das A, Bandopadhyay B, Jha TB (2012) *In vitro* propagation and molecular evaluation of *Capsicum annuum* L. cultivar with high chromosome number (2n=48). Scientia Horticulturæ 140: 119-124.
- Salvador MH (2002) Genetic resources of chilli (*Capsicum* spp.) in Mexico. Proceedings of the 16th International Pepper Conference, Tampico, Tamaulipas, Mexico, pp: 10-12.
- Dhaliwal MS (2007) Solanaceous vegetables. Handbook of vegetable crops. Kalyani Publishers, Ludhiana, pp: 34-76.
- Purkayastha J, Alam SI, Gogoi HK, Singh L (2012) *Capsicum assamicum* sp. nov. (Solanaceae), from Assam, northeastern India. Ozean J Appl Sci 5: 55-66.
- Simonne AH, Simonne EH, Eitenmiller RR, Mills HA, Green NR (1997) Ascorbic acid and provitamin A contents in unusually colored bell peppers (*Capsicum annuum* L.). J Food Composition Anal 10: 299-311.
- Reyes-Escogido MDL, Gonzalez-Mondragon EG, Vazquez-Tzompantzi E (2011) Chemical and pharmacological aspects of capsaicin. Molecules 16: 1253-1270.
- Ravishankar GA, Suresh B, Giridhar P, Rao SR, Johnson TS (2003) Biotechnological studies on Capsicum metabolite production and plant improvement. In: De AK, editor. Capsicum: The genus Capsicum. London: CRC Press.
- http://www.fao.org/faostat/en/#rankings/countries_by_commodity/
- Indian Horticulture Database (2011) National Horticulture Board. New Delhi: Aristo Printing Press, pp: 6-7.
- Dutta SK, Singh SB, Saha S, Akoijam RS, Boopathi T, et al. (2017a) Diversity in bird's eye chilli (*Capsicum frutescens* L.) landraces of north-east India in terms of antioxidant activities. Proc Natl Acad Sci India Sect B Biol Sci 87: 1317.
- Mohammadi SA, Prasanna BM (2003) Analysis of genetic diversity in crop plants - Salient statistical tools and considerations. Crop Sci 43: 1235-1248.
- Bosland PW, Baral JB (2007) BhutJolokia - The world's hottest known chilli pepper is a putative naturally occurring interspecific hybrid. Horticult Sci 42: 222-224.
- Bhagowati RR, Changkija S (2009) Genetic variability and traditional practices in Naga King Chili landraces of Nagaland. Asian Agri History 13: 171-180.
- Dubey RK, Singh V, Upadhyay G, Pandey AK (2015) Assessment of phytochemical composition and antioxidant potential in some indigenous chilli genotypes from northeast India. Food Chem 188: 119-125.
- Sanatombi K, Sen-Mandi S, Sharma GJ (2010) DNA profiling of Capsicum landraces of Manipur. Scientia Horticulturæ 124: 405-408.
- Yumnam JS, Tyagi W, Pandey A, Ng TM, Rai M (2012) Evaluation of genetic diversity of chilli landraces from north eastern India based on morphology, SSR markers and the Pun1 Locus. Plant Mol Biol Rep 30: 1470-1479.
- Rai RP, Glint VD, Babu KN (2013) *In vitro* plant regeneration in *Capsicum chinense* Jacq. (Naga Chili). J Appl Biol Biotechnol 3: 30-33.
- Yatung T, Dubey RK, Singh V, Upadhyay G (2014a) Genetic diversity of chilli (*Capsicum annuum* L.) genotypes of India based on morphochemical traits. Aust J Crop Sci 8: 97-102.
- Yatung T, Dubey RK, Singh V, Upadhyay G, Singh S (2014b) Studies on seed protein profiling in chilli (*Capsicum annuum* L.) genotypes of northeast India. Aust J Crop Sci 8: 369-377.
- Kundu S, Das A, Haque SM, Ghosh B (2015) Chemotypic diversity in different Bhut Jolokia fruits: *In vitro* conservation and mass propagation of superior ecotype. Int J Pharm Sci Rev Res 34: 47-53.
- Dutta S, Singh SB, Vanlalhmangaiha, Banerjee A, Akoijam RS, et al. (2017b) *Capsicum frutescens* L. landraces of northeast India: from phenotypic diversity perspective of unexplored collection. Proc Natl Acad Sci India Sect B Biol Sci, pp: 1-12.
- Alice AK, Dubey RK, Pandey AK, Singh V, Singh S (2017) Studies on genetic diversity in chilli (*Capsicum annuum* L.) through SDS-PAGE protein profiling. Int J Chem Stud 5: 465-470.
- Adluri PK, Borah AR, Nath PD (2017a) Study of genetic diversity of Bhut Jolokia germplasm in north east India by SSR markers and morphology. Int J Pure Appl Biosci 5: 1657-1665.
- Bharathi LK, Rengasamy S, Singh R, Prabhu KV, Sharma A, et al. (2011). Estimation of capsaicin and

- capsaicinoid contents of high pungent chilli accessions of Andaman and Nicobar Islands and northeast India. Indian J Horticult 68: 551-555.
28. Kehie M, Kumaria S, Tandon P (2013) *In vitro* plantlet regeneration from cotyledon segments of *Capsicum chinense* Jacq. cv. Naga King Chilli and determination of capsaicin content in fruits of *in vitro* propagated plants by high performance liquid chromatography. Scientia Horticulturæ 164: 1-8.
 29. Sweat KG, Broatch J, Borrer C, Hagan K, Cahill TM (2016) Variability in capsaicinoid content and scoville heat ratings of commercially grown Jalapeño, Habanero and Bhut Jolokia peppers. Food Chem 210: 606-612.
 30. Ananthan R, Subhash K, Longvah T (2018) Capsaicinoids, amino acid and fatty acid profiles in different fruit components of the world hottest Naga king chilli (*Capsicum chinense* Jacq.). Food Chem 238: 51-57.
 31. Moirangthem SS, Gogoi S, Thongbam PD, Ramya KT, Fiyaz RA, et al. (2014). Effect of sowing time and crop geometry on the Capsaicinoid content in Bhoot Jolokia (*Capsicum chinense* Jacq.). Food Sci Technol 51: 1974-1981
 32. Bhutia KL, Ng TM, Khanna VK (2018) *In vitro* direct regeneration of Dalle Khursani (*Capsicum annum*) from salicylic acid treated explants. J Pharmacogn Phytochem 7: 1008-1012.
 33. Sarpras M, Gaur R, Sharma V, Chhapekar SS, Das J, et al. (2016) Comparative analysis of fruit metabolites and pungency candidate genes expression between Bhut Jolokia and other *Capsicum* species. PLoS One 11: e0167791.
 34. Meghvansi MK, Siddiqui S, Khan MH, Gupta VK, Vairale MG, et al. (2010). Naga chilli: A potential source of capsaicinoids with broad-spectrum ethnopharmacological applications. J Ethnopharmacol 132: 1-14.
 35. Liu Y, Nair MG (2010) Capsaicinoids in the hottest pepper Bhut Jolokia and its antioxidant and anti-inflammatory activities. Nat Prod Commun 5: 91-94.
 36. Baruah S, Zaman MK, Rajbongshi P, Das S (2014) A review on recent researches on Bhut Jolokia and pharmacological activity of capsaicin. Int J Pharm Sci Rev Res 24: 89-94.
 37. Venkataiah P, Subhash K (2003) Research articles: Thidiazuron induced high frequency adventitious shoot formation and plant regeneration in *Capsicum annum* L. J Plant Biotechnol 5: 245-250.
 38. Pandey U, Pandey J (2002) Anti-bacterial properties of cyanobacteria: A cost-effective and eco-friendly approach to control bacterial leaf spot disease of chilli. Curr Sci 82: 262-264.
 39. Garg R, Kumar S, Kumar R, Loganathan M, Saha S, et al. (2013). Novel source of resistance and differential reactions on chilli fruit infected by *Colletotrichum capsici*. Australasian Plant Pathol 42: 227.
 40. Adluri PK, Baldoldiya GM, Nath PD (2017b) Screening of Bhut Jolokia (*Capsicum chinense* Jacq.) germplasm of northeast India against chilli leaf curl virus. Int J Pure Appl Biosci 5: 1189-1196.
 41. Bhutia ND, Seth T, Shende VD, Dutta S, Chattopadhyay A (2015) Estimation of heterosis, dominance effect and genetic control of fresh fruit yield, quality and leaf curl disease severity traits of chilli pepper (*Capsicum annum* L.). Scientia Horticulturæ 182: 47-55.
 42. Rajapakshe RGAS, Ranasinghe JADAR (2002) Development of variety screening method for anthracnose disease of chilli (*Capsicum annum* L.) under field conditions. Trop Agric Res Extension 5: 7-11.
 43. Kumar R, Barman A, Jha G, Ray S (2013) Identification and establishment of genomic identity of *Ralstonia solanacearum* isolated from a wilted chilli plant at Tezpur, northeast India. Curr Sci 105: 1571-1578.
 44. Banerjee A, Dutta R, Roy S, Ngachan SV (2014) First report of chilli veinal mottle virus in Naga chilli (*Capsicum chinense*) in Meghalaya, India. Virus Dis 25: 142-143.
 45. Pandey KK, Gupta RC (2015) Management of anthracnose (*Colletotrichum capsici*) in chilli (*Capsicum annum* L.) through fungicides, bioagents and hand picking methods. J Spices Aromatic Crops 24: 141-144.
 46. Ng TC, Singh YH, Sumitra PH, Singh S, Singh SR, et al. (2017). Molecular based indexing of viral disease complex of king chilli (*Capsicum chinense* J.) in north eastern region of India. J Pharmacogn Phytochem 6: 2004-2008.
 47. Sanatombi K, Sharma GJ (2006) *In vitro* regeneration and mass multiplication of *Capsicum annum* L. J Food Agric Environ 4: 205-208.
 48. Sanatombi K, Sharma GJ (2007). Micropropagation of *Capsicum frutescens* L. using axillary shoot explants. Scientia Horticulturæ 113: 96-99.
 49. Sanatombi K, Sharma GJ (2008a) *In vitro* propagation of *Capsicum chinense* Jacq. Biologia Plantarum 52: 517-520.

50. Sanatombi K, Sharma GJ (2008b) *In vitro* plant regeneration in six cultivars of *Capsicum* spp. using different explants. *Biologia Plantarum* 52: 141-145.
51. Sanatombi K, Sharma GJ (2012) *In vitro* regeneration of *Capsicum chinense* Jacq. *Curr Trends Biotechnol Pharm* 6: 66-72.
52. Kehie M, Kumaria S, Tandon P (2012) Osmotic stress induced-capsaicin production in suspension cultures of *Capsicum chinense* Jacq. cv. Naga king chili. *Acta Physiol Plant* 34: 2039.
53. Gogoi S, Acharjee S, Devi J (2014) *In vitro* plantlet regeneration of *Capsicum chinense* Jacq. cv. 'Bhut jalakia': Hottest chili of northeastern India. *In Vitro Cell Dev Biol Plant* 50: 235.
54. Bora G, Gogoi HK, Handique PJ (2014) Effect of silver nitrate and gibberellic acid on *in vitro* regeneration, flower induction and fruit development in Naga chilli. *Asia Pac J Mol Biol Biotechnol* 22: 137-144.
55. Mangang RKJ (2014) *In vitro* callus induction of placental tissues of *Capsicum chinense* Jacq. cv. 'umorok' using different concentrations and combinations of growth hormones. *Int J Interdiscip Multidiscip Stud* 1: 63-66.
56. Gayathri N, Gopalakrishnan M, Sekar T (2015) *In vitro* micropropagation of *Capsicum chinense* Jacq. (Naga king chili). *Asian J Plant Sci Res* 5: 13-18.
57. Bhutia KL, Ng TM, Khanna VK (2016) *In vitro* regeneration of Dalle Khursani, an important chilli cultivar of Sikkim, using various explants. *Agrotechnology* 5: 142.
58. Singh A, Singh RK, Sureja A (2007) Cultural significance and diversities of ethnic foods of northeast India. *Indian Journal of Traditional Knowledge* 6: 79-94.
59. Horticulture at a Glance (2017) Government of India, Ministry of Agriculture and Farmers Welfare, Horticulture Statistics Division
60. Sharma A (2016) Sustainable economic analysis and constraints faced by the Naga king chilli growers in Nagaland. *Indian J Agric Res* 50: 220-225.
61. <http://www.indianspices.com/html/s0623chl.htm>