# Journal of Agriculture and Forest Meteorology Research

JAFMR, 2(6): 237-245 www.scitcentral.com



**Original Research Article: Open Access** 

# Systemic plant pathogen *Botrytis cinerea* Association with Aphids *Myzus persicae* Influence the Growth of Host Plant

Yahaya SM<sup>\*</sup>, Mardiyya AY, Safiyanu I, Sakina SB, Sale AI and Hassan MI

\*Department of Biology, Kano University of Science and Technology, Wudil, P. M. B. 3244 Kano, Nigeria.

Received March 22, 2019; Accepted April 02, 2019; Published November 10, 2019

# ABSTRACT

Plant pathogens and insect herbivores attaching and surviving on plant and their produce can associate indirectly with each other and interact directly or indirectly resulting in serious economic loss to the growers. In this experiment the influence of co-interaction between systemic plant pathogen *Botrytis cinerea* Pears Fr (Helotiales Sclerotiniaceae) and herbivorous insect green peach aphid *Myzus persicae* on the growth of lettuce *Lactuca sativa* (Asteraceae: Compositae) was investigated. Studies showed that the co- interaction between *Botrytis cinerea* and *Myzus persicae* causes stress on the lettuce plants resulting in economic loss. The studies found that co-interaction between *B. cinerea* and *M. persicae* resulted in lower lesion of *B. cinerea* and *M. persicae* counts when both were present on same host plants. In addition the stress resulting from co-interaction affected the growth of lettuce plant resulting in a significant reduction in plant height, internode length, leaf size and also significant reduction of fresh shoot and root weight.

Keywords: Botrytis cinerea, Co-interaction, Plant growth, Myzus persicae

# INTRODUCTION

*Myzus persicae* and *Botrytis cinerea* an herbivorous insect and plant pathogenic fungus respectively are of great economic importance due to their ability to spread diseases causing loss of plants and its produce [1-3]. Therefore, the ability of *Botrytis cinerea* and *Myzus persicae* to survive on the same plant host makes them to exhibit a co-interaction.

*Myzus persicae* Sulzer (Hemiptera: Aphididae), the common green peach potato aphid, is found throughout the world, where it is principally regarded as a serious pest of many important agricultural crops because of its ability to transmit plant viruses [4]. It feeds on crops such as peaches, potatoes, sugar beet, tobacco and various ornamental crops grown in landscapes and in glasshouses and most lettuce plants [4,5]. A high population of *M. persicae* (Figure 1) on a crop causes injury by removing large volumes of sap from the plants and depleting them of needed nutrients [4]. In addition, they also cause indirect injury by the production of sugary honeydew which makes the leaves susceptible to microbial attack, which then reduces leaf quality [6].

*Botrytis cinerea* is a systemic plant pathogen with a wide host range of more than 500 plants [7]. It is the causal agent of grey-mold disease and causes greater economic losses, during both pre and post-harvest phase of plant growth, more than any other disease [2,8], it is also responsible for about 20% losses of affected crops worldwide [2]. It is reported to cause annual crop losses worldwide ranging from \$10-\$100 billion [7]. Plant and their produce stored for weeks or months at temperatures ranging from 0-10°C and transported fruit, small fruits, ornamental flowers and bulbs, as well as forest seedlings are all attacked and destroyed by B. cinerea [9-11]. Therefore, Williamson et al. [12] concluded that B. cinerea as the most widely distributed disease of vegetables, ornamental fruits and field crops throughout the world. The pathogen produces clear or grey conidia on branched conidiophores which are dispersed by humid air currents, splashing water, tools and clothing; the conidiophores initiate a new infection on healthy plants [2,13]. The conidia of the fungus can infect the seedlings, flowers, stems or leaves through wounded or senescing tissues and directly through the epidermis [2,14]. The symptoms of the pathogen may sometimes appear very

**Corresponding author**: Yahaya SM, Department of Biology, Kano University of Science and Technology, Wudil, P. M. B. 3244 Kano, Nigeria, E-mail: sanimyahya@yahaoo.com

**Citation:** Yahaya SM, Mardiyya AY, Safiyanu I, Sakina SB, Sale AI, et al. (2019) Systemic plant pathogen *Botrytis cinerea* Association with Aphids *Myzus persicae* Influence the Growth of Host Plant. J Agric Forest Meteorol Res, 2(6): 237-245.

**Copyright:** ©2019 Yahaya SM, Mardiyya AY, Safiyanu I, Sakina SB, Sale AI, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

quickly or an infection may remain quiescent and may appear later when the tissues age or during storage [15-17]. The fungus overwinters either as mycelium in decaying plant debris or as sclerotia which are considered to be the main survival structure of *B. cinerea* in the soil [12].

The lettuce Lactuca sativa L. (Asteraceae (Compositae)) is a significant horticultural crop, widely used as source of food [18-20]. Many researchers have reported studies of Botrytis cinerea on crops such as strawberries, kiwifruits, raspberries, grapes and others [15]. Lettuces seem to have received little attention with respect to their infection by B. cinerea and yet they share the devastating effect caused by B. cinerea under favorable conditions, especially in protected crops. The aphids affecting lettuce crops are a problem worldwide causing serious economic losses [21,22]. Higher number of aphids may result in stunted plant growth, but, the most damaging effect of aphid attack is wilting and head contamination, which reduce the market value of the lettuce [21]. Despite the fact that lettuce is attack by several species of aphids however, the most common aphid pest of lettuce is the green peach-potato aphid (Myzus persicae Sulzer) [4,21].

The co-interactional interaction of insect herbivores and pathogenic fungi inhabiting the same host plant occur in a situation where the first attacker alters the fitness of the plant host in a way that it affects the second attacker, while in return the second attacker induces the plant host to synthesised chemicals which reduces the growth of the first attacker all at the expense of other important plant processes [23-26]. However, the success of these types of interaction is dependent on the type and behavior of insect herbivore and fungal pathogen involved [27,28]. Therefore two hypotheses were tested which are: (i) Decrease in the spread and expression of B. cinerea and lower growth rate of aphids will occur following stress on the host lettuce plant by the co-interaction of systemic pathogen B. cinerea and aphids M. persicae. (ii) Reduction of plant height and dry mass of the plant will occur as a result of stress on the plant host due to co-interaction of systemic plant pathogens and aphids.

# MATERIALS AND METHODS

# **Study plants**

Sterilized seed of lettuce was sown in 80, 15 cm diameter pots filled with a vermiculite growing medium in a controlled environment room (18-20°C, ambient humidity and 12-14 h L: D). Half of the plants (40, plants) were grown from clean uninfected seeds while the remaining half were grown from seed systemically infected with *B. cinerea*.

# Plant infestation with the Myzus persicae

One month after germination all the experimental plants were infested with ten nymphs of aphids *Myzus persicae* Sulzer (Hemiptera Aphididae). The *M. persicae* were reared for three generation for proper effect of telescoping of

generation before being used in the experiment [29]. Infestation was done by placing the aphids on the reverse side of the leaves (20 infected and 20 uninfected plants) using moist brush. In order to prevent the escape of the aphids immediately after infestation plants were covered with a vented plastic container. The remaining uninfested plants (20 infected and 20 uninfected) served as controls.

# Population size of Myzus persicae

The population size of *M. persicae* was obtained by counting the number of aphids on the plants. The counting was done once a week for twenty weeks, starting four weeks after infestation. Also visual examination was used to assess the appearance of *B. cinerea* infection on the plants.

#### Plant height

Plant height was taken from all 80 plants in the four treatments. Height of the plants was measured before harvest using measuring tape.

# Leaf size

Leaf size was taken from all 80 plants in the four treatments. Before harvest leaf size was measured from the two fully expanded middle leaves using measuring tape.

#### Internode length

Before harvest Internode length was taken from all the 80 plants in the four treatments. Before harvest length of the internode was measured from all the plants by using measuring tape.

# Dry shoot weight (g)

Harvested shoots were removed from all the plants and washed under running tap water and allowed to dry in an oven for one hour. Thereafter, dry shoot weight was taken from all the plants using an electronic balance (Kern scale Technic, 440-21N).

#### Dry root weight (g)

The roots were washed under running tap water and allowed to dry in an oven for 1 h. The measurements were taken using an electronic balance (Kern scale Technic, 440-21N).

# STATISTICAL ANALYSIS

All data were analysed using ANOVA with post-hoc Tukey tests [30]. As the data from plant height, internode length, leaf size, fresh shoot and root weight did not meet assumptions of normality, a Box-Cox approach was used to determine the correct transformation prior to analysis. The data from plant height and internode length were log transformed while data from leaf size and fresh shoot weight was square root transformed before the analysis. Root weight was Ln transformed. Number of aphid colony survivorship between infected and uninfected plants was analysed using two ways ANOVA [31].

# RESULTS

# Expression of systemic *B. cinerea* lesion on *M. persicae* infested and uninfected plants

More lesion of *B. cinerea* was recorded on plants uninfected by aphids than on infested plants. However, uninfected plants were free of lesions. Therefore, the presence of *B. cinerea* significantly reduces the plant height, leaf size, internode length and plant dry weight (Table 1).

# Number of *Myzus persicae* on infected and uninfected plants by systemic *B. cinerea*

The increase in the number of *M. persicae* colonies was very slow on both infected and uninfected plants. Therefore, the

**Table 1.** Influence of *B. cinerea* and aphid infestation on plant height, leaf size, internode length and dry weight of lettuce plants.

Parameters	Plant treatment	Test statistics
Plant height	B. cinerea infection	F1,38=55.63, P<0.001
	Aphid infestation	F1,38=53.54, P<0.001
	Interaction term	F1,38=4.57, P<0.361
Leaf size	B. cinerea infection	F1,39=52.53, P<0.001
	Aphid infestation	F1,38=45.46, P<0.001
	Interaction term	F1,38=51.12, P<0.641
Internode length	B. cinerea infection	F1,38=33.43, P<0.001
	Aphid infestation	F1, 38=44.51, P<0.039
	Interaction term	F1, 38=0.15, P=0.476
Dry shoot weight	B. cinerea infection	F1,38=3.21, P<0.001
	Aphid infestation	F1,38=43.51, P<0.001
	Interaction term	F1,38=0.13, P<0.431
Dry root weight	B. cinerea infection	F1,39=63.52, P<0.001
	Aphid infestation	F1,38=52.63, P<0.001
	Interaction term	F1,38=34.38, P<0.001

#### Size of plant parts

**Influence of systemic** *B. cinerea* and *M. persicae* on plant height: There was a significant effect of aphid infestation status and *B. cinerea* infection status on the height of the plant was obtained from the experimental plants in which the presence of either resulted in a significant reduction in plant height. The interaction term was not significant (Figure 1).

239

rate of survival of *M. persicae* colonies was not affected significantly by plant infection status ( $F_{1,19}=0.43$ , P=0.363). Higher number of *M. persicae* was counted on uninfected plants, while the lowest count was recorded on infected plants. The effect of aphid infestation significantly decreases the plant height, leaf size, internode length and fresh weight of shoot and root of lettuce plants (**Table 1**).



Figure 1. Plant height (mm) for experimental plants in the presence and/or absence of B. cinerea and M. persicae.

# Knowledge of students about lawns in the campus environment

Influence of systemic *B. cinerea* and *M. persicae* on leaf size: There was a significant effect of aphid infestation

status and *B. cinerea* infection status on the leaf size of the experimental plants (Table 1) where the presence of either resulted in a significant reduction in leaf size. The interaction term was not significant (Figure 2).



Figure 2. Leaf size (mm) for experimental plants in the presence and/or absence of B. cinerea and M. persicae.

**Influence of systemic** *B. cinerea* and *M. persicae* on internode length: There was a significant effect of aphid infestation status and level of *B. cinerea* infection on the internode length of the experimental plants (Table 1) where the presence of either resulted in a significant reduction in leaf size. The interaction term was not significant (Figure 3).



Figure 3. Internode length (mm) for experimental plants in the presence and/or absence of B. cinerea and M. persicae.

# Plant dry weight

**Influence of systemic** *B. cinerea* and *M. persicae* on dry **shoot weight:** A significant effect of aphid infestation status and *B. cinerea* infection status on the dry shoot weight of the

experimental plants (Table 1) where the presence of either resulted in a significant reduction in dry shoot weight. However, the interaction term was not significant (Figure 4).



Figure 4. Mass (g) of shoots for experimental plants in the presence and/or absence of B. cinerea and M. persicae.

**Influence of systemic** *B. cinerea* and *M. persicae* on dry **root weight:** *Myzus persicae* infestation or infection by *B. cinerea* courses a significant decrease in the dry root weight

of the plants (Table 1). Because the primary influence of either natural enemy was not additive, the interaction was significant (Figure 5).



Figure 5. Mass of root (g) for experimental plants in the presence and/or absence of B. cinerea and M. persicae.

# DISCUSSION AND CONCLUSION

The result of the experiment showed that the dynamics of the two natural enemies have an influencing factor on one another. While lettuce plants infested with ten *M. persicae* died within ten weeks of initial infestation, but *B. cinerea* infection did not result in such mortality. Infection by *B. cinerea* and infestation by *M. persicae* reduced plant height, leaf size, internode length, plant dry weight. The rates of expression of *B. cinerea* lesions were however lower on aphid infested plants and aphid numbers were lower on plants infected by *B. cinerea*. In general it appears that the effect of aphid attack was greater than that of *B. cinerea* infection.

In a report by Anna et al. [32] showed that following aphid's infestation wastage of plants and their produce occur as a result of injection of phytotoxin during feeding, causing disintegration of chloroplast which decreases food production there by slowing rate of plant growth. Despite the fact that the mechanism by which aphids affect plant metabolism is not fully understood however, studies by Heil and Bostock [33], Swarbrick et al. [34] and Golawska et al. [35] reported that the induction of defence is costly, causing more need of assimilates by the plant. Also the ability of the herbivore to manipulate the plant's carbohydrate metabolism for its own use deprives the plant sufficient carbohydrate for metabolic activities [36]. From the result of the present study it is clear that plant height, internode length, leaf size and dry shoot and root weights were significantly reduced by the attack of M. persicae and B. cinerea. Although the interaction of aphid and *B. cinerea* significantly reduced the root weight of the lettuce plants, the interaction of aphid and B. cinerea did not significantly affect plant height, internode length, leaf size, shoot weight but their effect is exert independently of one another.

Despite the resource quality of the lettuce plant was not measured in this study. The result of the present study has found that the effects of systemic B. cinerea are detrimental to the increasing aphid population in lettuce plants. Infection by systemic B. cinerea which spreads into the plant tissues as the plant grows lowers the amount of nutrients available, making the plant less favourable for the attacking aphid. In a similar study on the effect of B. cinerea and aphids (Rhodobium porosum Sanderson) on rose plants (Rosa hybrida L.). Mouttet et al. [37] reported that B. cinerea induced the plant to synthesise secondary metabolites which either have toxic effects, aversive and/or anti-feedant effects on aphids which may subsequently attack. Such a negative relationship causes a reduction in the reproduction rate of M. *persicae*, thereby lowering its population size. This showed that co- interactions has the ability to change the ecological interactions and spatial distribution of the insect herbivore [37-41] in particular with regards to its interactions with its natural enemies.

Result of the study showed that expression of *B. cinerea* was drastically reduced after infestation with *M. persicae* when compared to non-infested plants. In a similar experiment Mouttet et al. [37] found that infestation with aphids (*Rhodobium porosum* Sanderson) on *B. cinerea* infected plant result in lower expression of *B. cinerea* and the infestation triggers the induction of Salicylic acid (SA)-dependent pathway around the infection site which kill cells of *B. cinerea* and stop it growth. But, continuous feeding by the aphids on the cell contents triggers the plant to induce the wound-response pathways, (JA) and (ET) dependent pathways which reduce the population of aphids [42-45].

In a related study by Delucchi [46] on the effect of M. *persicae* on the growth of its plant host, Brussels sprouts. A general reduction in dry and fresh weight, height, internode

length, leaf number and size was reported, where the most striking observation was reduction in dry root weight which was partly a due to the removal of assimilates, which were otherwise available for storage. Brussels sprouts which were subjected to probing and salivation by *M. persicae* and on which continued feeding by the aphid was prevented showed exactly a similar reduction accompanied by increased respiration of the plant.

Therefore, Delucchi [46] as well as Mackauer and Way [47] concluded that the increased respiration of the plant, a response to wounding and/or aphid salivation, contributed considerably to a reduction in plant height, leaf size and dry weight. However, finding by Heng-Moss et al. [48] reported that the reduction in the plant fresh and dry weight occurs due to the reduction of photosynthesis in leaves which have been injured by interaction of aphid infestation and pathogen attack due to an increased synthesis of defensive chemicals in response to the attack [49].

The present research has confirmed that co-interactional relationship between a systemic pathogen and an insect herbivore attacking lettuce plants reduces growth of the lettuce plant [50,51]. Therefore, the results of the experiment shows that B. cinerea affect aphid and also aphid affect B. cinerea and each of them stress the lettuce plant. In addition, the experiment confirmed the existence of a negative relationship between *M. persicae* and *B. cinerea* where they independently stress the host plant, and in addition to the reduction of the population growth rate of each other, they also reduce the growth rate of the host plant, probably by triggering the induction of defence chemicals by the plant at the expense of other important vital functions [52-55]. This shows that B. cinerea may have far-reaching effects on coexisting insect herbivores. Therefore this research will serve as a valuable pointer in increasing our understanding of the ecological consequences of a ubiquitous but hitherto understudied interaction [56,57].

#### REFERENCES

- 1. Mackean DG (2004) Aphids, Hemiptera, life cycle, behavior reproduction and feeding damage to plants and control of damage. Available at: http://www.biologyresources.com/aphid-01.html
- 2. Agrios NG (2005) Plant Pathology. Elsevier Academic Press, USA.
- 3. Yahaya SM, Ahmed I (2015) Does seed serve as inoculum source for systemic *Botrytis cinerea* infection in lettuce. India J Plant Sci 4: 7-21.
- 4. Blackman RL, Eastop VF (2000) Aphids on the world's herbaceous plants and shrubs. Wiley, Chichester, England.
- 5. Yahaya SM, Mardiyya AY (2019) Review of postharvest losses of fruits and vegetables. Biomed J Sci Tech Res 13.

- Michael JS, Jennifer ST, Bart PHJT (2005) Plant mediated interactions between pathogenic microorganisms and herbivorous arthropods. Annu Rev Entomol 51: 663-689.
- 7. Hua L, Yong C, Zhaquan Z, Boqiang L, Guozheng Q, et al. (2018) Pathogenic mechanism and control strategies of *Botrytis cinerea* causing post-harvest decay in fruits and vegetables. Food Qual Saf 3: 111-119.
- Samir D, Amnon L (2007) Post-harvest Botrytis infection: Etiology, development and management. In Elad Y, Botrytis: Biology, pathology and control. Springer, Dordrecht, the Netherlands, pp: 349-362.
- 9. Govrin EM, Levine A (2000) The hypersensitive response facilitates plant infection by the necrotrophic pathogen *Botrytis cinerea*. Curr Biol 10: 751-757.
- Elad Y, Stewart A (2004) Microbial control of Botrytis cinerea. In Elad Y, Botrytis: Biology, Pathology and Control. Springer, Dordrecht, the Netherlands, pp: 223-236.
- 11. Yahaya SM, Fagwalawa LD, Lawan M (2015) Influence of airborne and seed inoculum in the initiation of leaf, stem and root infection by systemic *Botrytis cinerea* in Lettuce (*Lactuca sativa*). J Plant Pathol Microbiol 11.
- Williamson B, Tudzynski B, Tudzynski P, Van Kan JAL (2007) *Botrytis cinerea*: The cause of grey mould diseases. Mol Plant Pathol 8: 561-580.
- 13. Holz G, Coertze S, Williamson B (2004) The ecology of Botrytis on plant surfaces. In Elad Y, Botrytis: Biology, pathology and control. Springer, Dordrecht, the Netherlands, pp: 9-27.
- 14. Van Kan JAL (2005) Infection strategies of *Botrytis cinerea*. Acta Horticulturae 669: 77-89.
- 15. Sowley ENK (2006) Epidemiology of *Botrytis cinerea*. PhD Thesis, University of Reading, UK.
- 16. Elias SNK, Shaw MW, Dewey FM (2010) Persistent symptomless, systemic and seed-borne infection of lettuce by *Botrytis cinerea*. Eur J Plant Pathol 126: 61-71.
- 17. Yahaya SM, Mardiyya AY, Sakina SB, Hayatu LW (2019) Disease cycle and infection strategies of systemic plant pathogen *Botrytis cinerea*. Novel Res Microbiol J 3: 204-214.
- Norman JC (1992) Tropical vegetable crops. Stockwell Ltd., New York, pp: 13-53.
- 19. Ramu S, Kuppusamy P, Lingan R, Gandi K (2011) Efficacy of different biological control agents against major post-harvest pathogens of grapes under room

temperature storage conditions. Phytopathology 50: 55-65.

- Yahaya SM, Fagwalawa LD, Ali MU, Umma M, Lawan M, et al. (2016) Symptomless transmission of seed-borne *Botrytis cinerea* into the seed of next generation lettuce plant (*Lactuca sativa*). J Plant Sci Res 1.
- Palumbo JC (2001) Influence of admire and platinum on the population growth of the lettuce aphid under field condition. In Byrne DN and Baciewicz P. (ed.) Vegetable Report Series. University of Arizona, College of Agriculture and Life Science, pp: 14-19.
- Nicolaos EM, Dimitris EG (2002) Fungal and bacterial disease. In Emerging Infectious Diseases. Kluwer Academic Publishers: New York, pp: 34-46.
- Barbosa P (1991) Plant pathogens and non-vector herbivores. In: Barbosa P, Krischik VA and Jones CG (ed.). Microbial Mediation of Plant-Herbivore Interactions. Wiley, New York, pp: 341-382.
- Hatcher PE, Paul ND, Ayres PG, Whittaker JB (1994b) The effect of foliar disease (rust) on the development of *Gastrophysa viridula* (Coleoptera: Chrysomelidae). Ecol Entomol 19: 349-360.
- 25. Stout MJ, Thaler JS, Thomma BPJ (2006) Plantmediated interactions between pathogenic microorganisms and herbivorous arthropods. Annu Rev Entomol 51: 668-689.
- 26. Rodger G, Rahier M, Naisbit RE (2007) Coping with an antagonist: the impact of a phytophatogenic fungus on the development and behavior of two alpine leaf beetles. Oikos 116: 1514-1523.
- 27. Kruess A (2002) Indirect interaction between a fungal plant pathogen and a herbivorous beetle of the weed *Cirsium arvense*. Oecologia 130: 563-569.
- Cory JS, Hoover K (2006) Plant-mediated effects in insect-pathogen interactions. Trends Ecol Evol 21: 278-286.
- 29. Dixon AFG (1985) Structure of aphid populations. Annu Rev Entomol 30: 155-174.
- Hilton A, Armstrong R (2006) Post-hoc Tukey ANOVA test. Available at: http://eprints.aston.ac.uk/93317/1/
- Rehman MK (2013) Problem solving and data analysis using Minitab: A clear and easy guide to six sigma methodology. Wiley, Available at: http://www.ebookee.org/Problem-Solving-and-Data-Analysis-9
- Anna MB, Leon V, Carlos SJ, Francois N, Burger V (2012) Phloem feeding insect stress and photosynthetic gene expression. In: Najafpour M. (ed.) Applied photosynthesis. Springer: USA, pp: 22-33.

- 33. Heil M, Bostock RM (2002) Induced systemic resistance (ISR) against pathogens in the context of induced plant defenses. Annu Bot 89: 503-512.
- 34. Swarbrick PJ, Schuze-Lefert P, Scholes JD (2006) Metabolic consequences of susceptibility and resistance (race-specific and broad-spectrum) in barley leaves challenged with powdery mildew. Plant Cell Environ 29: 1061-1076.
- 35. Golawska S, Kryzyzanowski R, Lukasik (2010) Relationship between aphid infestation and chlorophyll content in Fabaceae species. ACTA Biologica Botanica 52: 76-80.
- 36. Rodriguez-Soana C, Law BR, Rufus I (2011). Manipulation of natural enemies in agroecosystems habitat and semiochemicals for sustainable insect pest control. In Larramendy ML and Soloneski S (ed.). Integrated Pest Management and Control Current and Future Tactics. InTech, pp: 89-126.
- 37. Mouttet R, Philippe B, Thomas C, Desneux N (2011) Phytophagous arthropods and a pathogen sharing a host plant. PLoS One 6: e18840.
- 38. Paul ND, Hatcher PE, Taylor PE (2000) Coping with multiple enemies: An integration of molecular and ecological perspectives. Trends Plant Sci 5: 220-225.
- Ohgushi T (2005) Indirect interaction webs: herbivoreinduced effects through trait change in plants. In Ohgushi T, Schmitz O and Holt RD (ed.) Ecological Reviews. Trait-Mediated Indirect Interactions. Cambridge University Press, p: 183.
- 40. Viswanathan DV, McNickle G, Thaler JS (2008) Heterogeneity of plant phenotypes caused by herbivorespecific induced responses influences the spatial distribution of herbivores. Ecol Entomol 33: 86-94.
- 41. Mauck KE, De Morales CM, Mescher MC (2010) Effects of cucumber mosaic virus infection on vector and non-vector herbivores of squash. Communicative and Integrative Biology 3: 579-582.
- 42. Kanno H, Fujita Y (2003) Induced systemic resistance to rice blast fungus in rice plant infested by whitebacked plant hopper. Entomologia Experimentalis et Applicata 107: 155-158.
- De vos M, Van Zaanen W, Koornef A, Korzelius JP, Dicke M (2006) Herbivore-induced resistance against microbial pathogens in Arabidopsis. Plant Physiol 142: 352-363.
- 44. Hatano E, Kunert G, Michaud JP, Wolfgang W, Weisser W (2008) Chemical cues mediating aphid location by natural enemies. Eur J Entomol 105: 797-806.

- 45. Wassim EC, Roy K, Tatyana S, Heather R, Florence N, et al. (2008) Distinct roles of jasmonates and aldehydes in plant-defense response. PLoS One 3: e1904.
- Delucchi VL (1976) Studies in biological control. International Programme, Cambridge University Press 9: 414-420.
- Mackauer M, Way MJ (1976) Myzus persicae Sulzer an aphid of world importance. In Delucchi VL (ed.). Studies in Biological Control. University Press: Cambridge, UK, pp: 51-119.
- 48. Heng-Moss TM, Macedo T, Markwell JP, Baxebdale FP, Quisenberyy SS, et al. (2003) Comparison of chlorophyll and carotenoid concentrations among Russian wheat aphid (Homoptera: Aphididae) infested wheat isolines. J Econ Entomol 96: 475-481.
- 49. Blackman RL (1974) Life cycle variation of *Myzus persicae* (Sulzer) (Homoptera: Aphidiidae) in different parts of the world, in relation to genotype and environment. Bull Entomol Res 63: 595-609.
- 50. Dixon AFG (1998) Aphid ecology: An optimization approach. Chapman and Hall: London, pp: 2-26.
- Eastop VF (1986) Aphid plant associations. Coevolution and Systematics. Clarenden Press: Oxford, pp: 33-54.
- 52. James WC (1974) Assessment of plant diseases and losses. Ann Rev Phytopathol 12: 27-48.
- 53. Margaritopoulos JT, Tzortzi M, Zarpas KD, Tsitsipis JA, Blackman RL (2006) Morphological discrimination of *Aphis gossypii* (Hemiptera: Aphididae) populations feeding on Compositae. Bull Entomol Res 96: 153-165.
- 54. Minitab Inc. (2009) Minitab statistical software, Release 16 for Windows, State College, Pennsylvania.
- 55. O'Neil TM, Gladders P, Ann DM (1997a) Prospects for integrated control of lettuce diseases. In: Crop Protection and Food Quality Meeting Customer Needs. Proceedings of BCPC/ANPP Conference. University of Kent, UK, p: 485.
- 56. Palumbo JC, Mullis J, Reyes F, Amaya A (1998) New insecticide alternatives for aphid management in head lettuce. In Oebker NFE (ed) Vegetable Report Series. University of Arizona. College of Agriculture and Life Science, pp: 1-10.
- 57. Synder W, Ives A (2003) Interaction between specialist and generalist natural enemies: Parasitoids, predators, pea aphids biocontrol. Ecology 84: 91-97.